

Hypertension and heart failure with preserved ejection fraction: position paper by the European Society of Hypertension

Alexandros Kasiakogias^a, Enrico Agabiti Rosei^b, Miguel Camafort^c, Georg Ehret^d, Luca Faconti^e, João Pedro Ferreira^{f,g,h}, Jana Brguljanⁱ, Andrzej Januszewicz^j, Thomas Kahan^k, Athanasios Manolis^l, Konstantinos Tsioufis^a, Thomas Weber^m, Thomas G. von Luederⁿ, Otto A. Smiseth^o, Kristian Wachtell^p, Sverre E. Kjeldsen^q, Faiez Zannad^{f,g,h}, Giuseppe Mancina^r, and Reinhold Kreutz^s

Hypertension constitutes a major risk factor for heart failure with preserved ejection fraction (HFpEF). HFpEF is a prevalent clinical syndrome with increased cardiovascular morbidity and mortality. Specific guideline-directed medical therapy (GDMT) for HFpEF is not established due to lack of positive outcome data from randomized controlled trials (RCTs) and limitations of available studies. Although available evidence is limited, control of blood pressure (BP) is widely regarded as central to the prevention and clinical care in HFpEF. Thus, in current guidelines including the 2018 European Society of Cardiology (ESC) and European Society of Hypertension (ESH) Guidelines, blockade of the renin-angiotensin system (RAS) with either angiotensin-converting enzyme inhibitors or angiotensin receptor blockers provides the backbone of BP-lowering therapy in hypertensive patients. Although superiority of RAS blockers has not been clearly shown in dedicated RCTs designed for HFpEF, we propose that this core drug treatment strategy is also applicable for hypertensive patients with HFpEF with the addition of some modifications. The latter apply to the use of spironolactone apart from the treatment of resistant hypertension and the use of the angiotensin receptor neprilysin inhibitor. In addition, novel agents such as sodium-glucose co-transporter-2 inhibitors, currently already indicated for high-risk patients with diabetes to reduce heart failure hospitalizations, and finerenone represent promising therapies and results from ongoing RCTs are eagerly awaited. The development of an effective and practical classification of HFpEF phenotypes and GDMT through dedicated high-quality RCTs are major unmet needs in hypertension research and calls for action.

Keywords: heart failure, heart failure hospitalization, hypertension, preserved ejection fraction, guidelines

Abbreviations: ABPM, ambulatory blood pressure monitoring; ACEi, angiotensin-converting enzyme inhibitor; ARB, angiotensin receptor blocker; ARNi, angiotensin receptor neprilysin inhibitor; BNP, brain natriuretic peptide; BP, blood pressure; CCB, calcium channel blocker; CKD, chronic kidney disease; ESC, European Society of Cardiology; ESH, European Society of Hypertension; FDA, Food and Drug Administration; GDMT, guideline directed

medical therapy; HF, heart failure; HFpEF, HF with preserved ejection fraction; HFREF, HF with reduced ejection fraction; HMOD, hypertension mediated organ damage; HR, hazard ratio; LV, left ventricle, left ventricular; LVEF, left ventricular ejection fraction; LVH, left ventricular hypertrophy; MRA, mineralocorticoid-receptor antagonist; NYHA, New York Heart Association; RAS, Renin-angiotensin system; RCT, randomized controlled trial; RR, rate ratio; SGLT2, sodium-glucose co-transporter-2

INTRODUCTION

Heart failure (HF) represents a major and highly relevant clinical consequence of hypertension-mediated organ damage (HMOD) [1]. Up to half of all patients presenting with HF have HF with preserved

Journal of Hypertension 2021, 39:1522–1545

^aFirst Department of Cardiology, Hippokraton Hospital, Medical School, National and Kapodistrian University of Athens, Athens, Greece, ^bDepartment of Clinical and Experimental Sciences, University of Brescia, Brescia, Italy, ^cInternal Medicine Department, ICMID, Hospital Clinic, University of Barcelona, Barcelona, Spain, ^dCardiology, Geneva University Hospitals, Geneva, Switzerland, ^eKing's College London British Heart Foundation Centre, London, UK, ^fUniversité de Lorraine, Centre d'Investigations Cliniques Plurithématique Inserm 1433, ^gCHRU de Nancy, Inserm U1116, ^hFCRIN INI-CRCT, Nancy, France, ⁱUniversity Medical Centre Ljubljana, Hypertension Department, Medical University Ljubljana, Ljubljana, Slovenia, ^jDepartment of Hypertension, National Institute of Cardiology, Warsaw, Poland, ^kKarolinska Institutet, Department of Clinical Sciences, Danderyd Hospital, Division of Cardiovascular Medicine, Stockholm, Sweden, ^lCardiology Department, Asklepeion General Hospital, Athens, Greece, ^mCardiology Department, Klinikum Wels-Grieskirchen, Wels, Austria, ⁿØkern Heart Centre, ^oInstitute for Surgical Research and Department of Cardiology, Oslo University Hospital, Rikshospitalet, and University of Oslo, ^pDepartment of Cardiology, Oslo University Hospital, ^qInstitute of Clinical Medicine, Medical Faculty, University of Oslo, and Departments of Cardiology and Nephrology, Oslo University Hospital, Ullevaal, Oslo, Norway, ^rUniversità Milano-Bicocca, Milan; Policlinico di Monza, Monza, Italy and ^sCharité - Universitätsmedizin Berlin, Humboldt-Universität zu Berlin, and Berlin Institute of Health, Institut für Klinische Pharmakologie und Toxikologie, Berlin, Germany

Correspondence to Reinhold Kreutz, Institute of Clinical Pharmacology and Toxicology, Charitéplatz 1, D-10117 Berlin, Germany. Tel: +49 30 450 525 112; e-mail: reinhold.kreutz@charite.de

Received 3 March 2021 **Revised** 27 April 2021 **Accepted** 8 May 2021

J Hypertens 39:1522–1545 Copyright © 2021 Wolters Kluwer Health, Inc. All rights reserved.

DOI:10.1097/HJH.0000000000002910

ejection fraction (HFpEF) and increasing evidence suggests that their risk of death and recurrent hospitalization is similar to patients with HF with reduced ejection fraction (HFrEF) [2]. The prevalence of hypertension among patients with HFpEF ranges between 55 and 90% and patients with HFpEF are more likely to present with a history of hypertension compared with those with HFrEF [3,4]. In a patient with an established diagnosis of HFpEF, hypertension, when present, may be the only cause or one of multiple causes or comorbidities. Even though available evidence is limited, appropriate control of blood pressure (BP) is widely regarded as central to the prevention and clinical care of HFpEF patients [5–7].

In the 2018 European Society of Cardiology (ESC) and European Society of Hypertension (ESH) Guidelines for the management of hypertension, a simplified core drug treatment algorithm is recommended for most hypertensive patients, including patients with uncomplicated hypertension, HMOD, diabetes and the elderly [1]. Guideline-directed medical therapy (GDMT) includes a combination of a blocker of the renin-angiotensin system (RAS), that is an angiotensin-converting enzyme inhibitor (ACEi) or angiotensin receptor blocker (ARB), with a calcium channel blocker (CCB) or thiazide/thiazide-like diuretic as initial therapy for most patients [1]. In addition, the steroidal mineralocorticoid-receptor-antagonist (MRA) spironolactone is recommended as the preferred drug for treating patients with resistant hypertension, that is patients in whom BP control is not achieved with a triple combination of a RAS blocker, CCB and thiazide/thiazide-like diuretic [1,8]. Even though there are no specific pharmacologic options for HFpEF, based on the current guideline recommendations, a RAS blocker would also represent the backbone of therapy in hypertensive patients with HFpEF, while treatment with an MRA would apply only in case of resistant hypertension. Nevertheless, although the core algorithm of GDMT for hypertension appears appropriate for patients with HMOD including patients with left ventricular (LV) hypertrophy (LVH), its applicability for the treatment of HFpEF patients could be critically questioned [1].

Clear evidence for medical treatments that improve the course of HFpEF represents one of the major unmet needs in cardiovascular medicine, which applies to an already large and increasing number of patients, most of whom are hypertensive. Thus, the optimal treatment strategy for hypertensive patients with concomitant HFpEF is currently also unknown due to the paucity of positive outcome data derived from randomized controlled trials (RCTs) and methodological limitations of the available studies [9–11]. Importantly, there is no definite evidence for any drug treatment, including the use of BP-lowering drugs, to support clear improvements in cardiovascular outcomes and mortality in HFpEF patients [11]. Nevertheless, targeting the RAS is fundamental to the BP-lowering algorithm, as it is the cornerstone of pharmacological treatment across the cardiovascular continuum with a significant body of evidence supporting the use of RAS blockers in hypertensive patients [1,11,12]. However, definitive superiority for the use of a RAS blocker (ACEi or ARB) or an MRA (spironolactone) has not been shown in RCTs explicitly designed for HFpEF [10,11].

The current position paper by the ESH seeks to review some of the pathophysiological aspects of HFpEF of particular interest to hypertensive patients, re-examine medical options investigated in major HFpEF trials and propose a current framework for treatment and future research directions for patients with hypertension and HFpEF. In this regard, the BP-lowering therapy of hypertensive patients with concomitant HFpEF but also the options for management of HFpEF beyond BP control are discussed. It should be acknowledged that although the approach to control BP in hypertensive patients with HFpEF could be considered to be distinct from that of treating HFpEF *per se*, current drug choices for therapy of these two conditions are largely overlapping. Nevertheless, the proposed recommendations in the current document would primarily apply to patients with HFpEF and a preceding or new diagnosis of hypertension.

PATHOPHYSIOLOGY OF HEART FAILURE WITH PRESERVED EJECTION FRACTION IN HYPERTENSION: IMPLICATIONS FOR TREATMENT

HFpEF is a heterogeneous condition in terms of pathophysiological pathways and clinical presentation [9,13]. It has been suggested that different etiologies contribute to a common pathophysiological substrate but also that assigning patients to more specific clinical sub-phenotypes may allow the identification and development of more targeted treatment strategies [9,14–17]. Similarly, in the context of hypertension, the variable contribution of diverse pathophysiological processes (e.g. type of geometric remodeling, degree of diastolic dysfunction, extent of myocardial fibrosis, neurohormonal activation and microvascular disease) may serve to produce different HFpEF phenotypes, which may require adjustments in the approach to treatment. In addition, even though it remains unclear whether HFpEF is a distinct entity or an intermediate step from normal to reduced LV systolic function, the transition from preserved to reduced LV ejection fraction (LVEF) in hypertensive patients is commonly considered the result of poorly controlled long-term sustained hypertension or incident myocardial infarction [18,19]. Overall, the pathophysiology of HFpEF in hypertensive patients is complex and multifactorial (Fig. 1), but certain vital issues merit consideration as they may have implications for management.

Left ventricular hypertrophy and the limitations of the left ventricular ejection fraction

The presence of LVH and/or LV diastolic dysfunction are considered cardinal features to establish the diagnosis of HFpEF, although they are not pathognomonic [5,6,20,21]. Echocardiographic data from large studies have shown that LVH may be absent in approximately half to three-fourths of patients with HFpEF [22,23]. Nevertheless, different types of remodelling patterns (including concentric and eccentric LVH and concentric remodelling) can be found in patients with hypertension with or without HFpEF [22–26]. The concentrically hypertrophied LV of the hypertensive patient, with preserved LVEF but with leftward/upward




FIGURE 1 Hypertension and HFpEF. The pathophysiologic milieu of hypertension (A) sets the stage for the development of further functional and structural changes in HFpEF (B) that underlie the clinical phenotype of symptomatic HFpEF patients (C). Selected mechanisms, risk factors and comorbidities are listed, while several other abnormalities not shown may be involved in HFpEF. Some mechanisms may contribute to more than one pillar across the disease continuum. Certain abnormalities may be evident only during exercise. The impairment of systolic and diastolic reserve and ventricular-arterial coupling, in conjunction with changes in preload, afterload, heart rate and rhythm, is central to the generation of symptoms. Comorbidities variably contribute to abnormalities throughout the disease process. cGMP, cyclic guanosine monophosphate; ECM, extracellular matrix; EF, ejection fraction; HFpEF, heart failure with preserved ejection fraction; LV, left ventricle; LVEDP, left ventricular end-diastolic pressure; NO, nitric oxide; RAS, renin-angiotensin-system.

shifting of its pressure volume-relationship reflecting increased filling pressures at rest and/or exercise has often served as an archetype of the pathophysiology of HFpEF, despite its limitations [9,13].

The LVEF is a simple but notoriously preload and afterload-dependent index that continues to serve as a convenient estimate of LV function, as well as for categorization of the failing heart, both in the clinical and research setting [27,28]. In clinical practice and RCTs, an arbitrary LVEF of less than 35 or 40% has been used to identify patients who have HF with reduced systolic function [5,6,27,28]. The use of a high LVEF cut-off of at least 50% compared to lower values, for example at least 40%, as a diagnostic criterion for HFpEF is recommended in recent guidelines [5]. However, a preserved LVEF does not exclude LV systolic function abnormalities, and, conversely, diastolic dysfunction often coexists with reduced LVEF [29,30].

The utility of the LVEF as a diagnostic tool for HFpEF is limited by the confounding effect of LVH. The LVEF is a parameter of chamber function and is used to express the

stroke volume as a fraction of end-diastolic volume. In concentric chamber remodelling, the LVEF may appear increased due to a reduction in LV cavity volume and not due to increased contractility [31]. In addition, there is a difference between LV chamber function (i.e. fractional shortening or LVEF) and myocardial function (i.e. midwall shortening) [32]. As the myocardial contractile elements are located in the midwall, the estimation of LV systolic function depends on the distance from the endocardium to the midwall. When performing measurements for estimation of LV chamber function, the wall is considered infinitely thin. However, this is a misleading assumption in the case of concentric remodelling and LVH. As a result, a seemingly normal LVEF could coexist with depressed midwall shortening, lower cardiac output and higher peripheral resistance [32–35]. Accordingly, LV longitudinal strain, that serves as a proxy for subendocardial function, appears to be a more meaningful measure compared to LVEF to describe alterations in contractility [36]. In hypertensive LVH, longitudinal LV strain may be affected despite an

apparently normal LVEF [37,38]. Impairments in longitudinal LV strain can be more pronounced in patients with HFpEF, and may have implications regarding prognosis and response to treatment [39].

Right heart involvement

The importance of right heart involvement in the clinical syndrome of HFpEF should not be overlooked [40,41]. Elevated left atrial pressures lead to pulmonary hypertension that may be further worsened by superimposed development of increased pulmonary vascular resistance and impaired pulmonary arterial compliance [42]. An elevated pulmonary artery systolic pressure (>35 mmHg) has been shown to effectively distinguish HFpEF patients from hypertensive individuals without HF [43]. Chronically increased pulmonary arterial pressures and the resultant right ventricular-pulmonary arterial coupling mismatch are main mediators of right ventricular dysfunction, observed in up to 50% of HFpEF patients and associated with a poor outcome [40,41,44]. Echocardiographic data in patients with a diagnosis of HFpEF and a previous history of pulmonary oedema or invasively confirmed increased filling pressures, showed that right ventricular systolic function and structure may deteriorate to a significantly greater extent over time when compared with the left ventricle (LV) [45].

Ventricular-arterial interaction

Using traditional pressure-volume loop analysis, patients with HFpEF show greater ventricular and arterial stiffening compared with what is expected from hypertension or ageing alone [46]. Their ratio may approximate its normal value and become less informative. Separate measurements of each component of the ratio with more sensitive markers of myocardial function (e.g. global longitudinal strain) and pulsatile arterial function (e.g. pulse wave velocity for arterial stiffness, augmentation index and reflection magnitude for wave reflections) have been recently recommended [47]. In particular, an increase in wave reflections may unfavourably increase late systolic load, which in turn impairs diastolic function [48].

Abnormalities in ventricular-arterial interaction in HFpEF may affect the clinical response and tolerance to commonly used BP-lowering drugs. Increased arterial stiffness, coupled with increased LV end-systolic elastance may lead to marked fluctuations in BP following changes in loading conditions or stroke volume [49]. The steep end-systolic pressure-volume relationships may thus lead to massive BP decreases without increasing stroke volume when vasodilators are used [50]. In addition, impaired chronotropic reserve along with reduced stroke volume reserve limit the expected increase in cardiac output during exercise [51]. This phenomenon may be further exacerbated with drug-induced heart rate lowering [52].

The effect of age

The majority of patients with HFpEF are over 65 years old and female. The syndrome of HFpEF is often regarded as an exaggerated presentation of the ageing heart. Older patients with HFpEF are more likely to be hypertensive, have lower body weight and exhibit more comorbidities as

well as a greater risk of underlying cardiac amyloidosis, compared with their younger peers [53–55]. Arterial and ventricular stiffening increase in parallel with age and in older hypertensive patients it is difficult to distinguish to what extent impaired LV diastolic parameters are due to ageing *per se* or are resulting from chronically elevated BP, that is increased afterload [56]. This poses diagnostic challenges and implications for risk assessment. Impairment of LV relaxation parameters in elderly with otherwise normal hearts is frequent and may be even prognostically benign. As a consequence, some authors have suggested to use age-based adjustments for indices of LV diastolic function [57]. Similarly, an increase in arterial stiffness and wave reflections, on top of the normal aging process, can be easily quantified and age and sex-specific reference values for central pulsatile haemodynamics are available [58].

Kidney disease and the cardiorenal syndrome

Hypertension is a major risk factor for chronic kidney disease (CKD), which is a common comorbidity in patients with HFpEF [1]. In turn, CKD has been associated with an increased risk of new-onset HFpEF and up to half of the patients with HFpEF have CKD defined as a glomerular filtration rate less than 60 ml/min per 1.73 m² [59]. There is evidence that CKD may be equally or even more strongly associated with mortality in patients with HFpEF than in those with HFrEF [60]. The association between CKD and HFpEF appears to be bidirectional [61]. The prevalence of LVH increases with worsening renal function [62], and patients with HFpEF and CKD have been suggested to have greater LV mass, more pronounced impairment of LV diastolic function and poorer ventricular and atrial strain measurements compared with individuals with preserved kidney function [63]. Proposed mechanisms implicated in the pathophysiological interactions between the heart and kidney in HFpEF (i.e. the cardiorenal syndrome) include increased central venous and intra-abdominal pressures, RAS activation, oxidative stress and chronic inflammation. It has been speculated that fibrosis could represent the unifying consequence of inflammation resulting from systemic diseases (such as hypertension or diabetes) and also promotes the various expressions of the cardiorenal syndrome continuum, including the HFpEF phenotype [64].

CRITICAL APPRAISAL OF CLINICAL STUDIES IN PATIENTS WITH HEART FAILURE WITH PRESERVED EJECTION FRACTION

Many well-designed landmark RCTs have provided robust evidence that ACEis, ARBs, beta-blockers and MRAs significantly improve mortality and morbidity in HFrEF [5,6]. A common denominator of these medications is their ability to produce neurohormonal blockade and suppress the activated RAS and sympathetic nervous system. Activation of the RAS and aldosterone pathway is an essential component in the pathophysiology of hypertension, LVH, cardiac remodelling and fibrosis, even though a direct causative contribution to HFpEF has not been demonstrated [9–12]. A number of trials have been conducted

in patients with HFpEF to investigate if the observed efficacy of relevant established treatments in hypertension and HFrfEF would extend to this patient group (Table 1) [65–71]. However, all these trials either failed or nearly missed to meet their primary endpoint, a finding that was surprising for most experts given the anticipated crucial role of the RAS in LVH and cardiac remodelling and the development of HF [7,11,12,72]. This has been also in contrast to the evidence of decreased mortality associated with the use of RAS blockers in observational studies [73,74].

It is recognized that the landmark RCTs deploying RAS blockade in HFpEF involved populations with heterogeneous characteristics and exhibited several weaknesses that may have underlined their failure as recently reviewed in detail (Table 2) [11]. It is important to note that in these studies BP was already overall well-controlled (<140/90 mmHg) at baseline, and there were substantial rates of concurrent treatment with other antihypertensive drugs. These factors may have impacted the effect size of the interventions [65–69,75,76]. It should also be considered that a substantial proportion of patients (e.g. elderly, people of Black ethnicity) do not respond or respond incompletely to RAS inhibition when used as first-line therapy or in combination drug therapy [77,78]. Finally, the extent of neurohormonal activation in HFpEF is less clear compared with HFrfEF [79,80], and while aldosterone associates with LVH in hypertensive patients, this may not be the case with left ventricular diastolic dysfunction [81]. Along these lines, a contemporary meta-analysis of the effects of neurohormonal inhibitors in HFpEF did identify significant benefit mostly driven by a reduction in HF hospitalizations [82]. In the following sections, a short overview of the design and limitations of major RCTs in HFpEF is presented.

TRIALS ON RENIN-ANGIOTENSIN SYSTEM BLOCKADE

Candesartan in heart failure: assessment of reduction in mortality and morbidity study (CHARM-PRESERVED)

In the CHARM program, HF subgrouping was performed before randomization with patients with LVEF of 40% or less allocated to CHARM-Added (candesartan vs. placebo added to standard treatment including ACEi) or CHARM-Alternative (if intolerant to ACEi) and patients with LVEF more than 40% allocated to CHARM-Preserved [83]. CHARM-Preserved included 3023 patients (65% with an LVEF \geq 50%) of whom two-thirds were hypertensive and almost half had experienced a myocardial infarction [65]. Ischemic heart disease as a cause of HFpEF was identified in 56% of patients. A reduction in the primary outcome (composite of cardiovascular death or first HF hospitalization) with candesartan compared with placebo failed to reach statistical significance; thus, the study was considered neutral. However, a trend for a reduction in HF hospitalizations was noted [hazard ratio 0.85, 95% confidence interval (95% CI) 0.72–1.01, $P=0.072$] and fewer patients were hospitalized with HF in the candesartan group.

Specific points require consideration when examining the conclusions of CHARM-Preserved. From a design

aspect, extensive nonrandomized use of beta-blockers (56% at baseline), CCBs (31%), ACEis (20%) and spironolactone (12%) and notable discontinuation rates of allocated study medication may have weakened the observed effect in the ARB group. Also, controlling for a large number of baseline covariates identified a more substantial adjusted hazard ratio (HR) of 0.86 (95% CI 0.74–1.00, $P=0.051$) for the primary composite endpoint and a HR of 0.84 (95% CI 0.70–1.00, $P=0.047$) for HF hospitalization. An analysis utilizing a method that takes into account repeat hospital admissions rather than only time to first event documented a significant reduction in the composite of recurrent HF hospitalizations and cardiovascular death [rate ratio (RR) 0.75, 95% CI 0.62–0.91, $P=0.003$] [84]. The overall positive result of the CHARM program for the combined endpoint with no evidence of heterogeneity by LVEF implies some benefit with candesartan for the entire LVEF range met in HF [85]. The results of CHARM-Preserved have led experts to give a weak recommendation for the use of ARB to reduce hospitalizations in HFpEF [6].

Perindopril in elderly people with chronic heart failure study (PEP-CHF)

In the PEP-CHF study, a total of 850 patients aged at least 70 years (79% hypertensive) diagnosed with HF due to diastolic dysfunction based on clinical and echocardiographic criteria were randomized to either perindopril 4 mg or placebo [66]. Treatment with diuretics as well as previous HF hospitalization were used to support the diagnosis further. No LVEF cut-off was set, but a LV wall motion index of less than 1.4 was used as an exclusion criterion serving as a proxy for reduced LVEF. All patients were followed until the last patient had completed at least one year of follow-up (median follow-up 2.1 years). There were an unexpected low event rate and a high rate of open-label ACEi use. After the 1-year visit, a significant number of patients stopped blinded treatment. At the end of the study, 35% of patients in the perindopril group and 37% assigned to placebo were on open-label ACEi. Thus, the steering committee decided to cease recruitment before reaching the initial target of 1000 participants due to the predicted lack of statistical power for analysis of the primary endpoint. The composite endpoint of total mortality and first HF hospitalization was not different between groups, with the obvious limitation that the low statistical power does not allow conclusions to be drawn. However, it was encouraging that within the first year, when most patients were on assigned therapy, treatment with perindopril was associated with improvements in symptoms and exercise capacity, and reductions in hospitalizations for HF (hazard ratio 0.63, 95% CI 0.41–0.97, $P=0.033$).

Irbesartan in heart failure with preserved ejection fraction study (I-PRESERVE)

I-PRESERVE studied the ARB irbesartan vs. placebo in 4133 patients with HFpEF (LVEF \geq 45%) who were aged at least 60 years and predominantly hypertensive (88%) [67]. Significant baseline symptoms and high rates of subsequent hospitalizations attested to an actual HF population. Hypertension was considered the primary cause of HFpEF in 64%

TABLE 1. Major randomized controlled trials in HFpEF

Trial, Year N Duration	Study drug Mean dose	Patient characteristics	NYHA	Previous HF hospitalization	Concurrent antihypertensive medications	Initial BP and HR (Means)	Decrease in BP and compared with placebo group	Endpoints and unadjusted outcomes
CHARM- Preserved [65], 2003 N = 3023 Median 37 months	Candesartan 25 mg	EF: 54% Age: 64 years Women: 40% BMI: 29 kg/m ² HT: 64% DM: 28%	Range: II-IV III-IV: 38%	69%	Diuretic: 75% β-blocker: 56% CCB: 31% ACE-I: 20% Spiro.: 11%	136/78 mmHg, 71 bpm	↓6.9 mmHg (SBP), ↓2.9 mmHg (DBP) at 6 months	CV death and first HF hospitalization: HR, 0.89; 95% CI 0.11-1.03; P=0.12 First HF hospitalization: HR, 0.85; 95% CI 0.72-1.01; P=0.072
PEP-CHF [66], 2006 N = 850 Mean 26 months	Perindopril 4 mg	EF: 64% Age: 76 years Women: 56% BMI: 28 kg/m ² HT: 79% DM: 21%	Range: I-IV III-IV: 24%	100%	Diuretic: 100% β-blocker: 55% CCB: 33% ACE-I: 36% Spiro.: 10%	139/80 mmHg, 73 bpm	Greater decrease in SBP and DBP (values not available)	All cause death & first HF hospitalization: HR, 0.92; 95% CI 0.70-1.21; P=0.55 First HF hospitalization: HR, 0.86; 95% CI 0.61-1.20; P=0.37
I-PRESERVE [67], 2008 N = 4128 Mean 49.5 months	Irbesartan 300 mg	EF: 59% Age: 72 years Women: 60% BMI: 30 kg/m ² HT: 88% DM: 27%	Range: II-IV III-IV: 80%	44%	Diuretic: 90% β-blocker: 73% CCB: 40% ACE-I: 40% Spiro.: 28%	136/79 mmHg, 71 bpm	↓3.6 mmHg (SBP), ↓1.9 mmHg (DBP) at 6 months	All cause death & first CV hospitalization: HR, 0.95; 95% CI 0.86-1.05; P=0.35 First CV hospitalization: HR, 0.95; 95% CI 0.85-1.08; P=0.44
TOPCAT [68,75], 2014 N = 3445 Mean 40 months	Spironolactone 25 mg	EF: 56% Age: 69 years Women: 52% BMI: 31 kg/m ² HT: 91% DM: 33%	Range: II-IV III-IV: 33%	72%	Diuretic: 82% β-blocker: 78% CCB: 37% ACE-I: 84% Spiro.: N/A	130/80 mmHg, 68 bpm	↓4.4 mmHg (SBP) (Americas) at 8 weeks	CV death & aborted cardiac arrest & first HF hospitalization: HR, 0.89; 95% CI 0.77-1.04; P=0.14 First HF hospitalization: HR, 0.83; 95% CI 0.69-0.99; P=0.04
PARAGON HF [69,76], 2019 N = 4796 Median 35 months	Sacubitril/ Valsartan 200 mg	EF: 58% Age: 73 years Women: 52% BMI: 30 kg/m ² HT: 95% DM: 43%	Range: II-IV III-IV: 20%	48%	Diuretic: 95% β-blocker: 80% CCB: 37% ACE-I: N/A Spiro.: 26%	131/74 mmHg, 70 bpm	↓4.5 mmHg (SBP) at 8 months	CV death & total HF hospitalization: RR, 0.87; 95% CI 0.75-1.01; P=0.06 Total HF hospitalization: RR, 0.85; 95% CI 0.72-1.00. P=0.056

ACE-I, angiotensin-converting enzyme inhibitor; BP, blood pressure; CCB, calcium channel blocker; CI, confidence intervals; CV, cardiovascular; DM, diabetes mellitus; EF, ejection fraction; HF, heart failure; HFpEF, heart failure with preserved ejection fraction; HR, heart rate or hazard ratio; HT, hypertension; NYHA, New York Heart Association; RR, rate ratio.

TABLE 2. Parameters that may have affected study results in major randomized controlled trials in HFpEF

Different accepted cut-offs of left ventricular ejection fraction for inclusion in the study
Markers of confirmation of HFpEF (e.g. indices of diastolic function, structural heart disease, elevated biomarkers, prior HF hospitalization)
Blood pressure control at study entry
Selection of placebo vs. active comparator for the control arm
Concurrent treatment with other antihypertensive/HF medication during follow-up
Selection of endpoints and composite outcomes (e.g. first vs. recurrent HF hospitalizations, urgent HF visits, total vs. cardiovascular mortality)
Other issues related to study conduct/data analysis (e.g. insufficient statistical power, early study termination, lack of adjustment for potential confounders, regional differences)

HFpEF, heart failure with preserved ejection fraction; HF, heart failure.

of cases and coronary artery disease in one-fourth. There was no difference in either the primary outcome (all-cause death and first cardiovascular hospitalization) or any of the secondary outcomes between groups.

There were significant limitations regarding the efficacy of irbesartan in patients with HFpEF. Use of ACEis was allowed if there was an indication other than hypertension, with a 33% cap in each participating centre. During the trial, use of nonrandomized medication reached 73% for beta-blocker, 40% for ACEi and 28% for spironolactone. The frequent nonrandomized use of RAS-blockers, MRA and beta-blockers may have produced a ceiling effect and prevented any added beneficial impact of irbesartan. In addition, a high rate of discontinuation of the study drug, reaching 34% by the end of the study, was observed. The primary composite outcome was all-cause death or hospitalization for a cardiovascular cause (HF, myocardial infarction, unstable angina, arrhythmia or stroke), making comparisons with other relevant studies less straightforward. Finally, adjustment for baseline differences was not performed. A recent exploratory analysis which adjusted for prognostic baseline variables routinely available in clinical practice, provided evidence of some significant benefit with irbesartan, documenting a hazard ratio of 0.89 (95% CI 0.80–0.99, $P=0.033$) for the primary composite and hazard ratio of 0.87 (95% CI 0.77–0.99, $P=0.039$) for the composite of cardiovascular death or HF hospitalization [86].

THE ROLE OF NEPRILYSIN INHIBITION COMBINED WITH ANGIOTENSIN RECEPTOR BLOCKADE

Neprilysin inhibition has been a tempting therapeutic target for HFpEF in hypertensive patients because the resultant augmentation in natriuretic peptides is expected to enhance vasodilation, increase diuresis/natriuresis, activate guanylyl cyclase, improve myocardial relaxation and reduce LV fibrosis and hypertrophy [87,88]. Stand-alone neprilysin inhibition also increases angiotensin II levels and therefore needs to be combined with a RAS blocker. From a mechanistic point of view, it has been recently shown that soluble neprilysin levels are lower in patients with HFpEF compared with healthy controls [89]. However, this does not necessarily reflect the respective neprilysin activity [89]. Of note, angiotensin receptor neprilysin inhibitors (ARNis) are very effective antihypertensive drugs, and studies in hypertensive patients showed greater reductions in BP as well as LV mass with ARNi compared with ARB treatment [90,91]. The first-in-class ARNi sacubitril/valsartan has been shown to decrease the risk of HF hospitalization

or cardiovascular death versus enalapril by 20% in patients with HFrEF [92]. A phase-II study documented greater reductions in NT-pro-brain natriuretic peptide (BNP), left atrial size, BP and dyspnoea with ARNi compared with valsartan in HF patients with LVEF more than 45% [93]. The Efficacy and Safety of LCZ696 Compared to Valsartan, on Morbidity and Mortality in Heart Failure Patients With Preserved Ejection Fraction trial (PARAGON-HF) was a randomized, double-blind study that examined the safety and efficacy of ARNi versus stand-alone valsartan on the composite outcome of total hospitalizations for HF (first and recurrent) and cardiovascular mortality in 4822 symptomatic patients aged 50 years or older with LVEF at least 45% (average 58%), LVH or left atrial enlargement, and increased natriuretic peptide levels [69]. Similar to patients in previous HFpEF trials, 96% of participants had hypertension, 43% had coronary artery disease, 43% had diabetes and 32% had atrial fibrillation.

In the PARAGON-HF trial, the reduction of the primary endpoint in the ARNi group narrowly failed to reach statistical significance (RR 0.87, 95% CI 0.75–1.01, $P=0.06$). There was, however, a marginally significant reduction in the number of total HF hospitalizations (RR 0.85, 95% CI 0.72–1.00, $P=0.056$). In addition, an expanded composite endpoint combining the primary endpoint with urgent HF visits was also significantly reduced in the ARNi group (RR 0.86, 95% CI 0.75–0.99, $P=0.04$) [94]. Prespecified subgroup analysis showed that patients with LVEF equal to or below the median (of 57%) as well as women derived a possible benefit (RR 0.78, 95% CI 0.64–0.95 and RR 0.73, 95% CI 0.59–0.90 for the primary endpoint, respectively) [69,95]. A prespecified pooled analysis combining clinical trial data on HFrEF and HFpEF populations showed that the greatest risk reduction with ARNi vs. RAS inhibition alone is observed at lower levels of LVEF, but this benefit extends to a higher LVEF in women compared to men [96]. A post-hoc analysis showed that initiation of treatment with the ARNi early after hospitalization for HF might be accompanied by a more pronounced risk reduction [97]. The ARNi group also presented with significant improvements in quality of life scores and New York Heart Association (NYHA) functional classification. Importantly, halving of the risk for a prespecified composite of renal events and worsening renal function (hazard ratio 0.50, 95% CI 0.33–0.77, $P=0.001$), with the treatment effect extending across the spectrum of baseline renal function, has been demonstrated [69,98]. In addition, combination of MRA with ARNi rather than valsartan appears to be associated with less decline in renal function [99]. It is important to underline that PARAGON-HF evaluated two distinct RAS blocking strategies, one with the

addition of neprilysin inhibition to valsartan, and the other with stand-alone valsartan as the active comparator. A putative placebo analysis of patient-level data from the major trials of ARNi and candesartan in heart failure patients suggested that there was indeed a clear treatment benefit of ARNi versus putative placebo for HF hospitalization or cardiovascular death across the full range of LVEF up to 60% [100].

The Randomized, Double-blind Controlled Study Comparing LCZ696 to Medical Therapy for Comorbidities in HFpEF Patients (PARALLAX, NCT number: NCT03066804) was recently presented in the ESC Congress 2020. The trial studied the effects of ARNi in 2566 NYHA II-IV patients (97% hypertensive) with LVEF more than 40% (mean 56%), evidence of structural heart disease and increased NT-pro-BNP [101]. The study population was stratified into three groups based on background therapy with RAS blockers (ACEi vs. ARB vs. no RAS blocker). With the use of such a three-arm parallel-group design, sacubitril/valsartan was tested against enalapril, valsartan or placebo. The ARNi versus standard medical therapy resulted in significant reductions in the surrogate marker of NT-pro-BNP at 12 weeks, but there was no additional benefit on 6-min walk distance, quality of life or NYHA class at 24 weeks [102]. A post-hoc analysis revealed an impressive 51% reduction in the exploratory endpoint of first HF hospitalizations at 24 weeks.

THE ROLE OF MINERALOCORTICOID RECEPTOR ANTAGONISTS

Treatment of Preserved Cardiac Function Heart Failure with an Aldosterone Antagonist trial (TOPCAT)

The use of MRA has been of particular interest for the management of HFpEF, considering their known benefits with respect to endothelial function and cardiac remodeling, particularly fibrosis and stiffness, but also arterial stiffness [103–105]. A positive signal had been observed in the small RCT Aldosterone Receptor Blockade in Diastolic Heart Failure (Aldo-DHF), which showed an improvement in echocardiographic markers of LV diastolic function with spironolactone [106]. TOPCAT investigated the prognostic effect of spironolactone versus placebo in 3445 predominantly hypertensive (91%) patients with HFpEF (LVEF \geq 45%) from the Americas (51%) and Russia/Georgia (49%) [68]. Patients were maintained on medical therapy with diuretics, beta-blockers and ACEis or ARBs. Spironolactone was uptitrated from 15 to 45 mg daily. The mean follow-up was 3.3 years, and the primary outcome was a composite endpoint of cardiovascular death, aborted cardiac arrest and HF hospitalization. The total event rate was 18.6 and 20.4% in the spironolactone and placebo groups, respectively, resulting in a nonsignificant trend for fewer events (hazard ratio 0.89, 95% CI 0.77–1.04, $P=0.14$) in favour of spironolactone.

Significant bias related to regional study conduct is strongly suspected of having influenced the actual trial results, even though the treatment-by-region interaction was not statistically significant. Firstly, patients in the Americas were more frequently enrolled based on increased BNP

levels compatible with HF, whereas investigators in Russia/Georgia largely enrolled patients based on recent HF hospitalization, a stratum raising the risk of an erroneous diagnosis of HF [68]. Secondly, there were marked regional variations in event rates, which were overall greater and significantly different between groups in the Americas (27.3% in the spironolactone vs. 31.8% in the placebo group; hazard ratio 0.82, 95% CI 0.69–0.98, $P=0.026$) compared with Russia/Georgia (9.3% in the spironolactone vs. 8.4% in the placebo group; hazard ratio 1.10, 95% CI 0.79–1.51, $P=0.576$) [107]. Thus, the placebo group in Russia/Georgia had a markedly lower occurrence of the primary outcome than in the Americas (8.4 vs. 31.8%), being as low as in a healthy population. Lastly, measurements of the serum concentration of the spironolactone metabolite canrenone in 366 patients who consented for the TOPCAT biorepository showed that an estimated 30% of individuals in Russia/Georgia randomized to spironolactone were not receiving or taking the drug [108].

Despite these shortcomings, in the entire population, treatment with spironolactone significantly reduced HF hospitalization rates (hazard ratio 0.83, 95% CI 0.69–0.99, $P=0.04$). In addition, in the Americas, spironolactone vs. placebo resulted in significantly lower event rates for the primary composite, cardiovascular death and HF hospitalization [107]. Most importantly, in the BNP stratification analysis, spironolactone resulted in lower rates in the primary composite endpoint (hazard ratio 0.65, 95% CI 0.49–0.87, $P=0.003$) [109]. Taken the available data together, one could consider that TOPCAT showed the superiority of spironolactone in the patients with biomarker-confirmed HFpEF who took their assigned medication. In addition, an analysis that examined outcomes across LVEF categories identified a greater estimated benefit of spironolactone at the lower end of the LVEF spectrum [110]. More light may be shed by the results of the ongoing Swedish Spironolactone Initiation Registry Randomized Interventional Trial in Heart Failure With Preserved Ejection Fraction (SPIRRIT, NCT02901184) and the German Spironolactone In the Treatment of Heart Failure Trial (SPIRIT-HF, EudraCT: 2017–000697-11). The ongoing Study to Evaluate the Efficacy and Safety of Finerenone on Morbidity and Mortality in Participants With Heart Failure and Left Ventricular Ejection Fraction Greater or Equal to 40% (FINEARTS-HF, NCT04435626) has been designed to evaluate the efficacy of the non-steroidal MRA finerenone to reduce cardiovascular death and HF events in patients with HF and an ejection fraction at least 40%.

THE ROLE OF BETA-BLOCKERS AND OTHER HEART RATE LOWERING DRUGS

Data from clinical trials suggested that higher heart rates are associated with worse outcomes in HFpEF patients in sinus rhythm [111]. From a physiological standpoint, impaired LV relaxation is associated with reducing LV stroke volume with increasing heart rates [9,13]. Beta-blockers have been traditionally considered of potential benefit because heart rate lowering would be expected to improve early diastolic filling of the stiff hypertrophied LV and reduce myocardial oxygen demand. However, certain caveats need consideration [52].

Heart rate slowing may only prolong diastasis without affecting LV filling at rest. Conversely, prolonged diastolic filling increases ventricular volumes and pressures, increasing the ventricular load and wall stress, as suggested by increased natriuretic peptides observed in patients on beta-blockers [112]. This mechanism may be detrimental for the predominantly older population of HFpEF patients with limited physical activity. In addition, heart rate slowing drugs may further aggravate chronotropic incompetence, and the limited chronotropic reserve may further impair exercise tolerance. There is a delicate balance between the increase in diastolic filling time and preservation of chronotropic reserve but at the expense of inefficient enhancement of relaxation with tachycardia [9,51,52]. Finally, the weaker central SBP reduction with beta-blockers compared with other antihypertensive drugs may provide an additional burden in this population [113].

Trial results on the effects of heart rate lowering in HFpEF patients have been mixed [114–116]. Of note, several relevant studies and meta-analyses have used the low LVEF cut-off of 40 or 45%, making the results difficult to interpret with respect to current HFpEF definitions [114–117]. In terms of intermediate endpoints, the randomized placebo-controlled Effects of Long-term Administration of Nebivolol on the clinical symptoms, exercise capacity, and left ventricular function of patients with Diastolic Dysfunction study (ELANDD) failed to show any benefit of nebivolol in symptoms or exercise capacity among 116 patients with an LVEF more than 45% and evidence of diastolic dysfunction, most of whom were hypertensive [118]. Even though the nitric oxide-releasing attributes of nebivolol may favourably affect aortic and ventricular compliance, the negative result was presumably associated with the concomitant inhibition of the heart rate response. Similarly, in HFpEF patients with a heart rate of at least 70 beats per minute, heart rate reduction with the *I_f* channel inhibitor ivabradine failed to improve filling pressures and functional capacity and reduce NT-pro-BNP levels [119].

With respect to hard endpoints, The Beta-blockers in Heart Failure Collaborative Group recently pooled individual patient-level data from double-blind RCTs in heart failure in order to examine the effects of beta-blockers on all-cause and cardiovascular mortality according to LVEF [120]. A total of 17312 patients from 11 major studies were included. Among patients in sinus rhythm at baseline, in the small group of 244 patients with HFpEF (LVEF > 50%), with a baseline median SBP of 147 mmHg, treatment with beta-blockers showed no benefit in terms of all-cause and cardiovascular morbidity. This was in contrast to the substantial benefit observed in the larger groups of HF with reduced LVEF (<40%) and mid-range HF (LVEF 40–49%), although the low number of trial patients included in this LVEF category represents a limitation. An explorative analysis of the American data of the TOPCAT trial showed that among 1567 mostly hypertensive participants with LVEF more than 50%, baseline beta-blocker use was associated with a greater risk of HF hospitalization (hazard ratio 1.74; 95% CI 1.28–2.37) [121]. In the same study, a higher LVEF was also associated with a greater risk among patients receiving beta-blockers, while a sensitivity analysis confirmed the results among patients who continued or discontinued beta-blocker therapy during follow-up [121].

POTENTIAL OF SODIUM-GLUCOSE CO-TRANSPORTER-2 INHIBITORS IN HEART FAILURE WITH PRESERVED EJECTION FRACTION PATIENTS WITH AND WITHOUT DIABETES

The sodium-glucose co-transporter-2 (SGLT2) is a low-affinity, high-capacity glucose transport protein located in the proximal convoluted tubule of the nephron and responsible for renal glucose reabsorption [122]. Several SGLT2 inhibitors are approved for treatment of type 2 diabetes as they have a glucose-lowering effect with a low risk of hypoglycaemia [123,124]. LV diastolic dysfunction is frequent in diabetes, and shows a strong correlation with insulin resistance and hyperglycaemia [125,126]. HFpEF represents the most frequent form of HF in patients with diabetes and is particularly prevalent in older female patients with hypertension [127]. SGLT2 inhibitors display multiple modes of action with particular interest to diabetic patients with hypertension and HFpEF as they not only reduce blood glucose levels, but also decrease BP, lower body weight and exhibit renal protection [128,129]. There is also evidence that SGLT2 inhibitors may improve volume regulation, reduce arterial stiffness, inhibit cardiac fibrosis, reduce LV mass, improve cardiac energetics and improve diastolic function [129–134].

Use of SGLT2 inhibitors has been shown to significantly lower the risk for hospital admissions for HF in type 2 diabetes mellitus patients at high cardiovascular risk, who were mostly hypertensive [135–138]. The risk reductions for HF hospitalizations were consistent and in the range of 27–39% [135–138]. Data from the Study to Evaluate the Effect of Dapagliflozin on the Incidence of Worsening Heart Failure or Cardiovascular Death in Patients With Chronic Heart Failure (DAPA-HF) and the Empagliflozin Outcome Trial in Patients With Chronic Heart Failure With Reduced Ejection Fraction (EMPEROR-Reduced) have supported the beneficial effect of SGLT2 inhibitors in HFrEF patients with and without type 2 diabetes [139,140]. A prespecified meta-analysis of these two single large-scale trials indicated consistent pooled treatment effects in HFrEF patients [141]. The benefit associated with SGLT2 inhibitors was primarily related to reducing HF hospitalizations, and, secondarily, to improved renal outcomes and decreased all-cause and cardiovascular death. Importantly, these benefits were seen regardless of age, sex, presence of diabetes and treatment with an ARNi [141].

Because of the consistent and convincing RCT data available for SGLT2 inhibitors, recent guidelines and position papers recommend SGLT2 inhibitors for patients with type 2 diabetes and established cardiovascular disease or multiple risk factors to prevent HF hospitalizations [123,124,128,142–145]. Similar recommendations extend to patients with HFrEF regardless of the presence of diabetes [144,145]. Regarding HFpEF, the ongoing Empagliflozin Outcome trial in Patients With Chronic Heart Failure With Preserved Ejection Fraction (EMPEROR-Preserved, NCT03057951) and Dapagliflozin Evaluation to Improve the Lives of Patients With Preserved Ejection Fraction Heart Failure trial (DELIVER, NCT03619213) are

of great interest as they are dedicated studies in HFpEF to assess the effect of empagliflozin and dapagliflozin on cardiovascular outcomes in patients with LVEF more than 40% with or without diabetes. Encouraging data were provided by the recently reported Sotagliflozin on Cardiovascular Events in Patients with Type 2 Diabetes Post Worsening Heart Failure (SOLOIST-WHF) RCT, which included patients with type 2 diabetes recently hospitalized for worsening HF (31% with an LVEF over 50%) [146]. Even though the trial was ended early because of loss of funding [147], it was shown that start of treatment with sotagliflozin around discharge of patients from hospital resulted in significantly lower risk for the primary end point of cardiovascular death and hospitalizations and urgent visits for HF. This treatment effect was consistent across subgroups of LVEF less than 50% or at least 50%.

OTHER TREATMENT OPTIONS

Many proof-of-concept studies, RCTs and observational analyses have examined other medical targets for HFpEF as recently reviewed [148,149]. Published results have either been disappointing (e.g. for organic nitrates) or requiring further investigation (statins, inorganic nitrates/nitrites, cytokine inhibitors, levosimendan). More trials examining novel treatments such as antifibrotic, anti-inflammatory and anti-oxidant agents, as well as cell therapies, are under way (Table 3). A detailed description of these trials falls outside the scope of this article, but specific points merit a mention. Phosphodiesterase-5 inhibitors have not been shown to improve exercise capacity in HFpEF patients [150], but may be beneficial for some patients with combined pre- and post-capillary pulmonary hypertension. This concept is currently investigated in the Phosphodiesterase-5 Inhibition in Patients With HF With Preserved Ejection Fraction and Combined Post- and Pre-Capillary Pulmonary Hypertension study (PASSION, EudraCT: 2017-003688-37).

Device-based therapies are also investigated and atrial unloading with a transcatheter interatrial shunt device has been shown to safely reduce capillary wedge pressures [151]. Catheter-based renal denervation has been shown to reduce LV mass and improve diastolic function [152], and recent randomized sham-controlled trials in hypertensive patients with or without treatment showed significant reductions in office and ambulatory BP [153,154]. Even though a small, underpowered trial in HFpEF patients failed to show a benefit with renal denervation [155], more trials are required to reveal the role of this method for hypertensive patients with HFpEF.

THERAPEUTIC APPROACH AND RECOMMENDATIONS FOR THE MANAGEMENT OF PATIENTS WITH HYPERTENSION AND HEART FAILURE WITH PRESERVED EJECTION FRACTION

In addition to BP control, the treatment of hypertensive patients with HFpEF seeks to minimize symptoms and improve functional capacity and quality of life, slow

progression of the disease, reduce the risk of HF hospitalizations and, ultimately, improve cardiovascular outcome and survival (Fig. 2). As a crucial step, vigorous management of concomitant risk factors and comorbidities is recommended to improve symptoms and/or prognosis even if relevant data is scarce [5,6,148,149,156]. Currently, no treatment has convincingly been shown to reduce mortality. Noteworthy, in clinical practice this predominantly elderly hypertensive population may be often frail, a characteristic that has been associated with increased dependency, more hospital admissions and greater mortality [157]. Therefore, secondary targets such as the reduction of hospitalizations and other patient-related outcomes, as, for example, improvements in quality-of-life metrics, are also important.

Regression of left ventricular hypertrophy and cardiovascular benefit

Treatment with all major antihypertensive drugs reduces LVH, although beta-blockers may be relatively less effective [72,158–161]. Several analyses from the Losartan Intervention for Endpoint reduction (LIFE) study have shown that in patients with hypertension, LVH regression induced by antihypertensive treatment was accompanied by improvements in indices of diastolic function and systolic performance [33,162]. In addition, reduction in LVH was associated with lower rates of clinical endpoints including new-onset HF, independently of BP reduction [7,163]. Although similar evidence regarding LVH regression in patients with HFpEF is not yet established, the study data in hypertensive patients with LVH support that LV mass reduction should also be pursued in individuals with HFpEF using strategies that lower cardiac afterload, peripheral vascular resistance and central BP [7,159,160].

Blood pressure targets

No study has directly investigated the optimal BP target in hypertensive patients with HFpEF [164]. Current recommendations may therefore only be based on extrapolations from populations with and without HF [1,164–167]. In the major randomized trials investigating RAS blockers and MRA, baseline BP was overall controlled and this may explain why further clinically non-negligible reductions in BP in the active compared with the control group (Table 1) were not accompanied by an identifiable decrease in cardiovascular endpoints [65–69]. For instance, in PARAGON-HF, the mean SBP at 8 months was lower by 4.5 mmHg in the ARNi arm compared with the valsartan group, a difference that was not associated with the treatment effect [69]. However, baseline and mean achieved SBP of 120–129 mmHg demonstrated the lowest risk for cardiovascular and renal outcomes [76]. An analysis of the TOP-CAT trial in 1645 participants from the Americas examined whether BP-lowering was associated with outcomes [75]. In the trial, the baseline BP was on average 126/71 mmHg and the authors identified a J-shaped association with adverse events as a SBP around 135 mmHg was associated with the lowest risk. A 4.4 mmHg reduction in SBP was observed in the spironolactone arm compared with the control group at 8 weeks. This reduction was similar across SBP quartiles

TABLE 3. Representative ongoing trials examining treatments in HFpEF

Trial name	Identifier ^a	Phase	Treatment	Sample size (Actual or estimated)	Follow-up for main endpoint	Main endpoint
Mineralocorticoid receptor antagonists						
FINEARTS-HF	NCT04435626	3	Finerenone	5500	42 months	Number of CV deaths or HF hospitalizations
SPIRIT-HFPEF	NCT02901184	3	Spironolactone	3200	5 years	CV death or first HF hospitalization
SPIRIT-HF	2017-000697-11	3	Spironolactone	1300	5 years	CV death or first HF hospitalization
Angiotensin receptor neprilysin inhibitors						
CNEPI	NCT03506412	4	Sacubitril/ Valsartan	40	5 weeks	Change in biomarkers that reflect neprilysin activity
PARAGLIDE-HF	NCT03988634	3	Sacubitril/ Valsartan	800	8 weeks	Proportional change in NT-pro-BNP
PERSPECTIVE	NCT02884206	3	Sacubitril/ Valsartan	592	3 years	Change in Global Cognitive Composite Score
PRISTINE HF	NCT04128891	3	Sacubitril/ Valsartan	60	12 months	Improvement in microvascular function and ischaemia
Beta-blockers						
CAYMUS HFpEF	NCT03948685	4	Carvedilol SR	300	24 weeks	Change in maximum NT-pro-BNP value
Preserve-HR	NCT03871803	4	Drug withdrawal	4	30 days	Change in peak VO ₂
Sodium-glucose co-transporter-2 inhibitors						
EMPEROR-Preserved	NCT03057951	3	Empagliflozin	5988	38 months	CV death or first HF hospitalization
EMPULSE	NCT04157751	3	Empagliflozin	500	90 days	Death or HF events or HF visits and change in KCCQ
DELIVER	NCT03619213	3	Dapagliflozin	6100	33 months	CV death rate or first HF hospitalization or urgent HF visit
DETERMINE- Preserved	NCT03877224	3	Dapagliflozin	504	16 weeks	Change in KCCQ and 6MWD
PRESERVED-HF	NCT03030235	4	Dapagliflozin	320	12 weeks	Change in KCCQ
Metformin						
PH-HFpEF	NCT03629340	2	Metformin	32	12 weeks	Mean pulmonary artery pressure during submaximal exercise
Inorganic nitrites/nitrates						
INABLE	NCT02713126	2	Sodium nitrite	100	12 weeks	Change in peak VO ₂
KNO3CK OUT HFPEF	NCT02840799	2	Potassium nitrate	76	12 weeks	Change in peak VO ₂ and total work during maximal-effort exercise test
ONOH	NCT02918552	2	Sodium nitrite	15	8 weeks	Change in peak VO ₂
PMED	NCT02980068	1	Oral nitrate	120	6 h	Change in nitrite/nitrate level in urine/plasma, bacterial content of oral/gut microbiome
Soluble guanylate cyclase stimulators						
DYNAMIC	NCT02744339	2	Riociguat	114	26 weeks	Change in cardiac output
Phosphodiesterase-5 inhibitors						
PASSION	2017-003688-37	2	Tadalafil	372	24 weeks	All-cause death or first HF hospitalization
Endothelin receptor antagonists						
SERENADE	NCT03153111	2	Macitentan	143	24 weeks	Change in NT-pro-BNP levels
Prostacyclin derivatives						
ILO-HOPE	NCT03620526	4	Iloprost	34	15 min	Change in pulmonary capillary wedge pressure after exercise
Antifibrotic agents						
PIROUETTE	NCT02932566	2	Pirfenidone	129	12 months	Change in extracellular volume in cardiac magnetic resonance
B3- Adrenergic receptor agonists						
BETA3_LVH	NCT02599480	2	Mirabegron	297	12 weeks	Change in left ventricular mass index, E/e'
Calcium sensitizer						
HELP	NCT03541603	2	Levosimendan	38	6 weeks	Change in PCWP
Fatty acid β-oxidation inhibitor						
DoPING-HFpEF	2018-002170-52	2	Trimetazidine	25	12 weeks	Change in PCWP
Anti-hyperuricemic drugs						
AMETHYST	2019-004862-16	2	Verinurad+ allopurinol	435	32 weeks	Change in exercise capacity
Cell therapy						
CELL-pEF	NCT02923609	2	CD34+ cell therapy	30	12 months	Change in E/e' assessed by cardiac magnetic resonance
Regress-HFpEF	NCT02941705	2	Allogeneic cardiosphere-derived cells	40	3 years	Safety profile during or post intracoronary delivery and during follow-up
Management of comorbidities/exercise therapy						
Cardiac Rehab Effects in HFpEF	NCT04506606	-	Afferent block	50	1 h	Femoral blood flow
FAIR-HFpEF	NCT03074591	2	Ferric carboxymaltose	200	12 months	Change in 6MWD
OPTIMIZE-HFPEF	NCT02425371	3	Management of comorbidities	410	2 years	Change in patient's well-being
PREFER-HF	NCT03833336	3	Ferric carboxymaltose	72	24 weeks	Change in 6MWD
Resistance Training in HFpEF	NCT02435667	-	Resistance Exercise Training	24	12 weeks	Bone density and strength, cardiopulmonary function, quality of life, blood biomarkers
Device therapies						
CCM-HFpEF	NCT03240237	Pilot	Cardiac contractility modulation	60	24 weeks	Change in KCCQ
CORolla	NCT02499601	-	CORolla TransApical Approach	10	6 months	All-cause mortality and serious adverse events
GUIDE-HF	NCT03387813	-	CardioMEMS	3600	12 months	Change in all-cause mortality, total number of HF hospitalizations, IV diuretic visits
PREFECTUS	NCT03338374	-	Cardiac resynchronization therapy	10	12 weeks	Change in diastolic and systolic reserve index in echocardiography
RAPID-HF	NCT02145351	-	Rate adaptive atrial pacing	30	4 weeks	Change in VO ₂ at ventilator anaerobic threshold
REDUCE LAP-HF II	NCT03088033	-	Interatrial shunt device	608	12 months	CV death or first non-fatal stroke or HF admissions and change in KCCQ

6MWD, 6-minute walk distance; CV, cardiovascular; HF, heart failure; HFpEF, heart failure with preserved ejection fraction; KCCQ, Kansas City cardiomyopathy questionnaire; NT-pro-BNP, NT-pro-brain natriuretic peptide; PCWP, pulmonary capillary wedge pressure.

^aIdentifier corresponds to clinical trial identifier in ClinicalTrials.gov or the European Union Clinical Trials Register.




FIGURE 2 Stepwise approach to the management of the hypertensive patient with HFpEF. BP, blood pressure; CKD, chronic kidney disease; COPD, chronic obstructive pulmonary disease; HFpEF, heart failure with preserved ejection fraction; OSA, obstructive sleep apnoea. *SBP targets should always be individualized, based on tolerability, particularly in frailer old or very old patients.

and overall sustained throughout the follow-up period, but with limited association with the observed risk reductions [75]. In an analysis of all TOPCAT participants with multiple available SBP measurements, a U-shaped association between a measure of mean SBP obtained during follow-up and mortality was observed, with SBP values of 120–129 and 130–139 mmHg associated with a lower risk [168]. Also, in a TOPCAT secondary data analysis restricted to patients enrolled in the Americas, spironolactone was associated with greater BP reductions (by 6.1 mmHg in SBP) and better BP control in HFpEF patients with characteristics of resistant hypertension. Importantly, the favourable effect of spironolactone on the primary outcome was similar in patients with and without resistant hypertension [169]. Finally, recent observational analyses from the Organized Program to Initiate Lifesaving Treatment in Hospitalized Patients with Heart Failure registry (OPTIMIZE-HF) provided some evidence that among older patients with HFpEF, a

discharge SBP less than 120 mmHg following hospitalization for HF may be associated with worse outcomes [170,171].

Indications for out-of-office BP measurements, namely ambulatory BP monitoring (ABPM) and home BP monitoring, include the diagnosis of white-coat, masked and resistant hypertension as well as the assessment of BP control and the night-time BP phenotypes [1]. However, published data on out-of-office BP monitoring in hypertensive patients with HFpEF is limited. Recent small observational cohort studies in Japanese patients hospitalized with HFpEF suggested that a riser pattern (a nocturnal BP fall <0% as assessed by ABPM) could be associated with all-cause mortality and cardiovascular outcome [172,173]. The Ambulatory Blood Pressure in HFpEF Outcomes Global Registry (HFPEFGlobal, NCT04065620) is an ongoing observational cohort study designed to assess the association of BP parameters derived from 24-h ABPM with cardiovascular

outcomes, taking also into considerations confounders such as comorbidities, frailty and functional capacity.

No properly designed RCT has been conducted to address BP treatment targets in HFpEF, but many major RCTs in hypertension have included patients with HMOD and HFpEF and have consistently shown that BP-lowering treatment effectively prevents HF events [1,167]. Hypertensive patients with HFpEF belong to the high/very high risk category, and, in agreement with the current recommendations of the 2018 ESC/ESH guidelines, prompt initiation of BP-lowering treatment alongside lifestyle interventions is indicated. In patients aged more than 80 years, even though the BP threshold for interventions to lower BP is generally at least 160/90 mmHg, antihypertensive treatment should also be considered when BP is at least 140/90 mmHg. In patients younger than 65 years, a target SBP in the range of 120–129 mmHg is recommended [1]; For older patients (age ≥ 65 years), a target SBP of 130–139 mmHg should be pursued, but treatment decisions should be individualized based on tolerability, frailty and comorbidities [1]. Accordingly, in older patients, initiation of combination therapy at the lowest available doses is recommended, while for the very old (>80 years), initiation of treatment with monotherapy may be appropriate. A target range between 70 and 79 mmHg for DBP applies to all patients with hypertension including those with HFpEF [1]. In line with current recommendations, it appears wise to avoid actively lowering BP to less than 120/70 mmHg, because the risk of harm may increase and outweigh the benefits [1,174].

Lifestyle changes

Even though there is limited evidence of the effect of different lifestyle measures in patients with HFpEF, promotion of lifestyle changes is recommended in order to improve BP control and possibly reduce cardiovascular events (Fig. 2). A large group of patients with HFpEF are obese and obesity is associated not only with insulin resistance and higher BP but also with reduced functional capacity and increased risk of developing HF [175]. A distinct phenotype of HFpEF patients with a BMI of at least 35 kg/m^2 showing more marked concentric LV remodeling, increased plasma volume and greater biventricular filling pressures with exercise has been proposed [176]. Although the optimal body weight target for hypertensive patients with HFpEF has not been established and an obesity paradox has been described in HF there is evidence of a beneficial effect of diet-induced weight loss on exercise capacity in obese patients with HFpEF [177]. Thus, the current ESC/ESH guideline 2018 recommendation for weight control to avoid obesity appears valid also for hypertensive patients with HFpEF [1].

The role of salt in the pathophysiology of fluid overload in HFpEF is unclear, and the evidence for salt restriction even for patients with HFpEF is weak [5,6,178]. In a small uncontrolled study of 13 hypertensive patients with HFpEF, a 3-week restricted sodium intake within the dietary approach to stop hypertension (DASH) diet led to improvements in LV diastolic function and arterial elastance [179]. Evidence for specific recommendations beyond the current guidelines [1] with respect to sodium intake in HFpEF

patients is lacking and there may be variations of optimal sodium intake based on symptoms, renal function and ethnic origin. Until more data are available, the current general recommendation of salt restriction to less than 5 g per day should be followed in all hypertensive patients, including the ones with HFpEF, particularly those prone to volume overload in order to reduce congestive symptoms [1]. Hypertensive patients with HFpEF should also be advised to pursue a balanced diet rich in vegetables, legumes, whole grains, fresh fruits, low-fat dairy products, fish and unsaturated fatty acids, and low in red meat and saturated fatty acids.

Regular exercise has been shown to improve cardiorespiratory fitness in patients with HFpEF and may as well affect prognosis. The Exercise Training in Diastolic Heart Failure pilot study (Ex-DHF) had shown that among 64 patients with HFpEF, a supervised combined endurance and resistance training program for three months was associated with improvements in peak $\dot{V}O_2$, markers of diastolic function and reported physical functioning [180]. A post-hoc analysis of the TOPCAT trial showed that HFpEF patients who reported ideal physical activity as per current recommendations (≥ 150 min/week of moderate aerobic activity or ≥ 75 min/week of vigorous activity) had lower rates of HF hospitalization and mortality compared with those with poor or intermediate level physical activity [181]. In an RCT which included 100 older obese individuals with HFpEF, 20 weeks of caloric restriction lead to a mean decrease of body weight by 7 kg, and a three-times-weekly supervised aerobic session individually and additively increased peak oxygen consumption [177]. Data from meta-analyses further support the concept that exercise training improves exercise capacity and quality of life, with less convincing evidence of a direct action on myocardial systolic or diastolic function [182]. However, very recently, the three-armed RCT Optimizing Exercise Training in Prevention and Treatment of Diastolic Heart Failure (OptimEx-Clin) performed on 180 patients with HFpEF (LVEF $> 50\%$) in NYHA class II or III documented no difference in peak oxygen consumption after 3 months of high-intensity interval vs. moderate continuous training, while neither mode met the prespecified difference in peak $\dot{V}O_2$ compared with one-time advice on physical activity according to guidelines [183].

Before prescribing an exercise program, baseline evaluation should include an exercise test (preferably cardiopulmonary exercise testing) in order to evaluate functional capacity, chronotropic reserve and BP response to exercise, monitor for ischemia or exercise-induced arrhythmia and accurately prescribe optimal exercise intensity. An exercise regimen tailored to the needs and capacity of each patient, which usually combines moderate-intensity endurance (3–5.9 metabolic equivalents) and dynamic exercise (30–50% of one-repetition maximum) is recommended [184]. Consideration should be made to include inspiratory muscle training (to enhance functional capacity) and strength training (to reduce sarcopenia) especially in certain high-risk groups such as elderly, frail patients with multiple comorbidities [185]. Implementation of exercise in the context of a cardiac rehabilitation program may be

considered depending on individual and resource settings, and patient preferences [184].

Drug therapy

The treatment goals in hypertensive patients with HFpEF are the control of BP *per se*, along with management of symptoms and improvement in prognosis. In Table 4, proposed recommendations for BP control in these patients and HFpEF-specific treatment are presented. It is suggested that, based on available data, the core drug treatment strategy of the 2018 ESC/ESH guidelines is also applicable for hypertensive patients with HFpEF (Fig. 3) [1]. The following modifications of the core treatment algorithm could be considered in HFpEF patients (Table 4, Fig. 3):

(1) **RAS blockers:** BP-lowering with a RAS blocker should be prescribed to all patients if not contraindicated. Apart from their BP-lowering action, available evidence supports a benefit in relation to LVH regression and diastolic function and has provided a positive signal for improvements in exercise capacity and decreases in HF

hospitalizations in HFpEF patients [65–67,70–74,159,162,163,186,187]. In addition, CKD and diabetes mellitus are common comorbidities in patients with HFpEF and RAS blockade has been well documented to reduce albuminuria and delay the progression of diabetic and non-diabetic CKD [1].

(2) **Diuretics:** The use of thiazide/thiazide-like diuretics is recommended in the current core treatment algorithm for BP control in hypertension, while loop diuretics are recommended in patients with glomerular filtration rates below 30 ml/min per 1.73 m². [1]. The thiazide-like diuretics chlorthalidone and indapamide, in particular, have been shown to efficiently reduce the risk of HF in elderly populations [1]. A differential and transiently intensified use of diuretics in HFpEF patients, most commonly loop diuretics, as compared to hypertensive patients with uncomplicated hypertension is often required for volume unloading in order to manage symptoms and avoid hospital visits [5,6,188,189]. A (transient) combination of loop diuretics with a thiazide diuretic, may be required in refractory cases. The optimal dosing of diuretics in HFpEF has not been

TABLE 4. Proposed recommendations for treatment of hypertensive patients with HFpEF

Recommendations ^a	Class	Level
Recommendations for BP control		
Prompt initiation of BP-lowering treatment is recommended simultaneously with lifestyle interventions when BP is $\geq 140/90$ mmHg [1].	I	A
Although initiation of BP-lowering treatment and lifestyle interventions in individuals aged >80 years is generally recommended when SBP is ≥ 160 mmHg, antihypertensive treatment should be considered in patients with HFpEF at this age when BP is $\geq 140/90$ mmHg and treatment is tolerated [1,5].	IIa	C
It is recommended that the first objective of treatment should be to lower BP to $<140/90$ mmHg and, provided that the treatment is well tolerated, treated BP values should be targeted to 130/80 mmHg or lower, in most patients [1].	I	A
In patients < 65 years, it is recommended that SBP should be lowered to a BP range of 120–129 mmHg [1].	I	A
In patients ≥ 65 years, it is recommended that SBP should be targeted to a BP range of 130–139 mmHg [1].	I	A
A DBP target in the range of 70–79 mmHg is recommended [1].	I	A
The BP threshold for initiation of treatment and the BP targets need to be individualized based on tolerability, frailty, and comorbidities, particularly for the elderly (>65 years). Close monitoring of adverse effects is recommended [1].	I	C
For BP control, combination treatment is recommended for most patients as initial therapy. Preferred combinations should comprise a RAS blocker (either an ACEi or an ARB) with a thiazide/thiazide-like diuretic or CCB [1].	I	A
It is recommended to initiate antihypertensive treatment with a two-drug combination, preferably in an SPC [1].	I	B
It is recommended that beta-blockers are combined with any of the other major drug classes for BP control, particularly when there are other specific clinical situations, e.g. angina, post-myocardial infarction or heart rate control [1,205].	I	A
For BP control, addition of low-dose spironolactone ^b , if tolerated, to existing BP-lowering therapy is recommended in patients with HFpEF and resistant hypertension [1,8,169].	I	B
HFpEF-specific recommendations		
Intensified use of diuretics, including loop diuretics, is recommended to relieve congestion and manage symptoms [5].	I	C
Irrespective from targeting BP, low-dose spironolactone ^b should be considered to reduce HF hospitalizations ^c [68,107,109].	IIa	B
Irrespective from targeting BP, use of sacubitril/valsartan (ARNi) as an alternative to ACE inhibitor or ARB should be considered to reduce worsening HF ^d [69,95–98].	IIa	B
SGLT2 inhibitors are recommended in hypertensive patients with diabetes mellitus to reduce the risk of HF hospitalization [128,135–138,142–146].	I	A

ACEi, angiotensin-converting enzyme inhibitor; ARB, angiotensin receptor blocker; ARNi, angiotensin receptor neprilysin inhibitor; BP, blood pressure; CCB, calcium channel blocker; HF, heart failure; HFpEF, heart failure with preserved ejection fraction; RAS, renin-angiotensin system; SGLT2, sodium-glucose co-transporter-2 inhibitor; SPC, single pill combination.

^aThe 2018 ESC/ESH guidelines did not include dedicated recommendations for HFpEF. Accordingly, the following recommendations are based on a critical appraisal of the general recommendations in these guidelines, novel data and reports with potential impact on HFpEF in hypertension.

^bThe dose range for spironolactone in the PATHWAY-2 study in resistant hypertension was 25 mg with forced titration to 50 mg once daily [8]. The dose range for spironolactone in the TOPCAT study in HFpEF was 15–45 mg once daily (mean dose 25 mg) [68]. A cautious use is recommended, particularly because of the risk of hyperkalaemia, in patients with an eGFR < 45 ml/min per 1.73 m² or baseline potassium ≥ 4.5 mmol/l based on PATHWAY-2 [8]. Patients with baseline potassium ≥ 5.0 mmol/l within the past 2 weeks or ≥ 5.5 mmol/l in the past 6 months, or with eGFR < 30 ml/min per 1.73 m² were excluded in TOPCAT [68].

^cA greater benefit of spironolactone was observed in patients with LVEF $< 50\%$ [68,110].

^dA greater benefit of sacubitril/valsartan was observed in patients with LVEF $\leq 57\%$ [69,96].




FIGURE 3 Drug treatment strategy for patients with hypertension and HFpEF. The core treatment algorithm for drug treatment of the ESC/ESH 2018 guidelines adapted for hypertensive patients with HFpEF. ACEi, angiotensin-converting enzyme inhibitor; ARB, angiotensin receptor blocker; ARNi, angiotensin receptor neprilysin inhibitor; CCB, calcium channel blocker; HFpEF, heart failure with preserved ejection fraction; MRA, mineralocorticoid receptor antagonist; SGLT2, sodium-glucose co-transporter-2.

established. Excessive use should be avoided considering that many patients may be preload-sensitive [190]. In patients with advanced symptoms, haemodynamic guidance with the use of a microsensor implanted in the pulmonary artery has been shown to significantly reduce hospitalizations [191]. Finally, the loop diuretic torsemide has been shown to improve diastolic function by positively affecting collagen cross-linking [192].

- (3) **MRA:** Spironolactone is the best studied MRA in hypertension as well as HFpEF. Although the use of low-dose spironolactone is recommended in patients with resistant hypertension [1,8,169], the addition of spironolactone should be considered earlier, and independently from the stage of BP control, to reduce hospitalizations for HF. Patients who may benefit more are those meeting the criteria of the TOPCAT trial (elevated BNP levels (BNP ≥ 100 pg/ml or N-terminal-pro-BNP ≥ 360 pg/ml, or hospital admission for HF within the previous 12 months), particularly those with a LVEF at the lower end of the spectrum [68,110]. Patients should be closely monitored for changes in potassium levels and renal function, and dosing of other diuretics should be adjusted based on clinical judgment (see Table 4 for further information regarding dosing). Very recently, the Food and Drug Administration (FDA) Cardiovascular and Renal Drugs Advisory Committee voted in favour of expanding the indication of spironolactone for the reduction of HF hospitalization in HFpEF patients [193,194]. A better understanding of the biological alterations in HFpEF and improved clinical phenotyping of HFpEF patients may provide more compelling evidence on the specific population that will mostly benefit from MRA [195,196]. Accordingly, the heart ‘Omics’ in AGEing RCT (HOMAGE) examined the

effects of spironolactone on people at increased risk of developing HF, who presented with features suggestive of clinically occult HFpEF. Spironolactone led to a significant decrease in SBP (by 10 mmHg) and NT-pro-BNP and improved echocardiographic markers of diastolic dysfunction, and, most importantly, appeared to favourably affect type-1 collagen turnover [196]. These findings are of particular interest, considering the role of myocardial fibrosis in patients with HFpEF and hypertension [197,198].

- (4) **ARNi:** For the time being, the results from PARAGON-HF, and the most recent PARALLAX studies have not provided unequivocal evidence on the superiority of ARNi over stand-alone ARB in HFpEF [69,102]. However, additional data and post-hoc analyses suggest that certain patient groups, such as women and those with a LVEF at the lower end of the HFpEF spectrum may derive benefit [69,95,96]. Therefore, ARNis should be considered as a replacement for conventional RAS blockers in these patient groups to reduce HF hospitalizations. It should not be dismissed that ARNis represent a very efficacious BP-lowering drug, which is, however, not approved for this indication [90,91,199]. Lastly, the same FDA committee as above, after evaluating the totality of available evidence voted also for an expanded indication for ARNi for HFpEF in the lower LVEF range [94,200]. Following this positive recommendation, the FDA has approved an indication for sacubitril/valsartan to reduce the risk of cardiovascular mortality and HF hospitalizations in adults with chronic HF [201]. Importantly, the new label states that benefits are most clearly evident in patients with LVEF below normal, but does not provide a LVEF cut-off, underlining the importance of clinical judgement in deciding who to treat [201].

- (5) **CCB:** In agreement with the general treatment algorithm, CCB appear as a valid choice in addition to RAS blocker therapy in patients with HFpEF, especially for younger patients or those free from clinical signs and symptoms of congestion. There is, however, limited data on the use of CCBs in HFpEF. Earlier small studies had shown improvements in diastolic function with non-dihydropyridine CCBs, but these results have not been examined in more recent trials [202]. The negative chronotropic and inotropic effect of non-dihydropyridine CCBs may be unsuitable for some patients, and regular follow-up to detect worsening LV function may be required. Non-dihydropyridine CCB are effective drugs for rate control in atrial fibrillation. The prognostic benefit of CCBs is unclear; in an analysis of 10 570 patients over 65 years old hospitalized with HFpEF (LVEF \geq 40%), a new discharge prescription for CCB was not associated with improvements in mortality, regardless of the class of the CCB [203,204].
- (6) **Beta-blockers:** In the 2018 ESC/ESH guidelines, beta-blockers are generally recommended at any treatment step when there is a specific indication in the core algorithm for patients with hypertension. However, although their use in HFrEF is clearly supported by a class I, level of evidence A indication, this does not apply to the HFpEF population [1,5,6,116]. On the contrary, there are even data supporting their cautious use in HFpEF patients, especially those with limited chronotropic reserve [51,52]. This contrasts with their frequent use in up to 80% of HFpEF patients as reflected in contemporary HFpEF studies (Table 1) [68,69]. Nevertheless, beta-blockers are effective antihypertensive agents and a very recent meta-analysis confirmed that treatment with beta-blockers results in significant reductions in cardiovascular endpoints, including HF events, that appear more pronounced among hypertensive patients [205]. In addition, an elevated heart rate has been associated with morbidity and mortality and a heart rate higher than 80 beats per minute has been included in the current ESC/ESH 2018 guidelines as a factor that influences cardiovascular risk [1,206]. Consequently, beta-blockers can be combined with any of the other major drug classes for BP control in HFpEF patients, especially those with other cardiac indications, for example angina, post-myocardial infarction or arrhythmias including atrial fibrillation requiring heart rate control.
- (7) **SGLT2 inhibitors:** Empagliflozin, canagliflozin, dapagliflozin, ertugliflozin and more recently sotagliflozin (a combined SGLT2 and SGLT1 inhibitor) have been consistently shown to reduce the risk of HF events in patients with diabetes mellitus and established cardiovascular disease or high cardiovascular risk [135–138]. They are therefore expected to be of significant benefit also for hypertensive patients with diabetes mellitus and HFpEF. When prescribing these medications, their BP-lowering and natriuretic effect should be taken into consideration. The results of upcoming RCTs shall provide

more evidence on the role of these drugs in patients with HFpEF with or without diabetes.

FUTURE DIRECTIONS

Phenotyping of heart failure with preserved ejection fraction

Patients with HFpEF present with remarkable diversity in terms of predisposing factors, pathophysiologic mechanisms, clinical patient profiles and level of noncardiac involvement [9,13–17,21]. Heterogeneity of HFpEF is already evident from the fact that not all HFpEF patients present with LVH or diastolic dysfunction [22–25]. Similarly, the fact that antifibrotic agents could be more effective in patients exhibiting a more pronounced fibrotic phenotype may partially explain why large trials of drugs targeting the RAS, which did not specifically include such patients, have failed to provide a solidly positive result [65–69,88,195–198,207–209]. Classification attempts based on epidemiological data, clinical characteristics, biological markers as well as machine-learning technology (phenomapping) have recently proven promising for identifying specific patient phenotypes who may benefit from more targeted treatment [14–17]. For instance, a combination of different clinical presentations and predisposing factors has provided twenty different HFpEF phenotypes that may be addressed in order to design individualized treatment [15]. The optimal approach in relation to phenotyping (e.g. pathophysiologic vs clinical) and how this may translate to personalized treatment and improve outcomes is a main area of scientific interest [14–17]. For example, lately there has been evidence that the burden of atrial fibrillation in patients with HFpEF may mark a distinct phenotype with impaired atrial mechanics, abnormal ventricular interactions and a different prognosis [210]. In terms of future research, it may appear more effective to investigate medical treatments in specific sub-populations (e.g. patients with/without obesity or atrial fibrillation, with phenotypes derived from machine learning-based clustering), rather than all-comers under the umbrella term of HFpEF [211].

Study methodologies and patient selection

A number of issues should be considered for future trials in patients with HFpEF. The taxonomy of HF has caused confusion with respect to interpretation of study results as well as treatment decisions. The current European definition of HFpEF requires an ejection fraction more than 50%, but a lower cut-off (e.g. $>45\%$ or $>40\%$) has been used in most landmark studies [5,6]. Post-hoc analyses of major trials have suggested a greater treatment effect in those patients with an LVEF at the lower end of the HFpEF range (e.g. LVEF $<50\%$), which has been labelled as mid-range HF [85,96,110]. Moreover, with speckle tracking echocardiography and measurement of longitudinal strain, it is possible to identify HFpEF with concurrent LV systolic dysfunction, which has been associated with a worse outcome [36–39,212]. The first phase ejection fraction, which corresponds to the percentage change in LV volume from end-diastole to peak ventricular fibre shortening, has also

TABLE 5. Upcoming advances and expectations related to the management of heart failure with preserved ejection fraction in hypertension

Establishment of most effective and practical classification of HFpEF patients (subphenotyping) to develop individualized and targeted therapies <ul style="list-style-type: none"> • Mechanistic/biological • Etiological • Clinical • Data-derived clustering
Optimization of study design to accurately include HFpEF patients <ul style="list-style-type: none"> • Utilization of a LVEF cut-off >50% • Consideration of subclinical systolic dysfunction • Utilization of undisputable markers of HFpEF (e.g. HF hospitalization, structural abnormalities, elevated biomarkers)
Determination of sex-related differences in disease presentation, prognosis and response to treatment
Determination of the role of atrial fibrillation in classification of HFpEF and disease course
Re-evaluation of established cardiovascular therapies (RAS blockade, MRA, ARNI) on specific HFpEF phenotypes, including patients with (uncontrolled) hypertension
Interpretation of the role of fibrosis and extracellular matrix homeostasis and the response to specific drug treatments
Identification of markers of greater response to certain pharmaceutical agents, with respect to symptoms and risk reduction
Trial results of promising therapies (SGLT2 inhibitors, sitagliptin, pirfenidone, mirabegron, finerenone, others)
Clarification of the contribution of lifestyle measures and exercise therapy to quality of life and prognosis

ARNI, angiotensin receptor neprilysin inhibitor; HF, heart failure; HFpEF, heart failure with preserved ejection fraction; LVEF, left ventricular ejection fraction; MRA = mineralocorticoid receptor antagonist; RAS, renin-angiotensin system; SGLT2, sodium-glucose co-transporter 2.

been shown to be impaired in hypertensive patients with diastolic dysfunction and could represent an early marker of systolic impairment despite a preserved LVEF [213]. Finally, following the recent approach of the FDA to the labelling of sacubitril/valsartan, where there was no mention of HF classification by LVEF [201], a new universal definition of HF may be contemplated [214–217]. This definition would be based on signs and symptoms, cardiac abnormalities and mode of presentation, further enriched by the cause of HF (e.g. hypertensive HF), rather than nominal LVEF values [214,215].

The addition of an elevated BNP or NT-pro-BNP as a biomarker criterion may further ensure that patients genuinely have HFpEF. However, normal levels observed in symptom-free periods and obese individuals may falsely preclude some patients from being enrolled in trials. At the same time, comorbidities (such as CKD or bouts of recurrent atrial fibrillation) or pharmacological interventions (such as beta-blockade) may increase circulating levels. The identification of markers of maximal clinical response to currently available or future treatments is an important objective in order to shape the optimal treatment for each patient. In this direction, analyses of I-PRESERVE and TOP-CAT have documented a greater prognostic benefit for irbesartan and spironolactone, respectively, in patients with lower BNP levels who may have less advanced disease [218,219]. However, no interaction has been shown between baseline NT-pro-BNP and the treatment effect of ARNi in PARAGON-HF [220].

CONCLUSION

We propose that further high-quality studies with careful patient selection fulfilling criteria of the dominant – in terms

of prevalence and cause – hypertensive HFpEF phenotype, will provide more definite evidence (Table 5). The studies need to be sufficiently powered and the study design should take into account the limitations as well as lessons learned from previous trials. Given the large volume of research demonstrating the effectiveness of RAS-blockers and MRA in hypertension, HFpEF and other cardiovascular diseases, these drug classes currently appear to have a potential in hypertensive patients with HFpEF not yet sufficiently explored. It remains to be seen whether the positive results of major hypertension trials that have investigated the aforementioned pillars of current cardiovascular therapy indeed extend to this group of patients. Finally, the results of already ongoing or planned studies using newer therapies such as SGLT2 inhibitors or the non-steroidal MRA finerenone in HFpEF patients may further modify the recommendations for the medical therapy of HFpEF *per se* and mainly in patients with hypertension due to the BP-lowering potential of these drugs.

ACKNOWLEDGEMENTS

Conflicts of interest

A.K. has no disclosures.

E.R.A. has received honoraria for lectures in the last 2 years from Menarini, Sanofi, Novartis, Pfizer, Eva Pharma, SunPharma, Hikma.

M.C. has received payment for advisory meetings from Novartis and from Boehringer Ingelheim.

G.E. has received payment for advisory/ speaker activities from Amgen, Bayer, Boehringer, Daiichi-Sankyo, Novartis, MSD, Servier and royalties from UpToDate.

L.F. has no disclosures.

J.P.F. is a consultant for Boehringer-Ingelheim, Astra Zeneca and Novartis.

J.B. has no disclosures.

A.J. has received personal fees and nonfinancial support from Servier, Krka, Berlin-Chemie/Menarini, Polpharma, and Gedeon Richter.

T.K. has research contracts between Karolinska Institutet and Medtronic, ReCor Medical and Amgen; all outside the presented work.

A.M. has received honoraria for lectures in symposia and educational activities sponsored by unrestricted grants from Amgen, Berlin-Chemie, Ferrer, Menarini, Sanofi-Aventis and Servier.

K.T. has received travel expenses or Research Grant or honoraria fees from the following: Medtronic, St. Jude Medical, Bayer, Novartis, Astra-Zeneca, Boehringer In, Pfizer, Chiesi, Pharmanel, Sanofi, Vianex, Win-Medica, ELPEN, Recordati, Servier.

T.W. has received honoraria for lectures and advisory board meetings in the last two years from Pfizer, Merck, Novartis, Servier, Boehringer Ingelheim, Daiichy Sankyo and Medtronic.

T.G.vL. has no disclosures.

O.S. has no disclosures.

K.W. has no disclosures.

S.E.K. has received Ad hoc lecture honoraria from Intas, Merck GBaA and Sanofi, and study committee honoraria from Takeda.

F.Z. reports steering committee personal fees from Applied Therapeutics, Amgen, Bayer, Boehringer, Novartis, Janssen, Cellprothera and CVRx, advisory board personal fees from, AstraZeneca, Vifor Fresenius, Cardior, Cereno pharmaceutical, Corvidia, Merck, Myokardia, NovoNordisk and Owkin, stock options at Cereno and G3Pharmaceutical, and being the founder of the Global Cardiovascular Clinical Trialist Forum.

G.M. has received honoraria for lectures and advisory board meetings in the last 24 months from Astra Zeneca, Boehringer Ingelheim, Daiichi Sankyo, Medtronic, Menarini, Merck, Novartis, Recordati, Sandoz, Sanofi, Servier.

R.K. has received support of research by Bayer and personal honoraria from Bayer, Berlin-Chemie Menarini, Daiichi Sankyo, Ferrer, Merck, Sanofi and Servier.

REFERENCES

- Williams B, Mancia G, Spiering W, Agabiti Rosei E, Azizi M, Burnier M, *et al.* 2018 ESC/ESH Guidelines for the management of arterial hypertension: The Task Force for the management of arterial hypertension of the European Society of Cardiology and the European Society of Hypertension. *J Hypertens* 2018; 36:1953–2041.
- Shah KS, Xu H, Matsouka RA, Bhatt DL, Heidenreich PA, Hernandez AF, *et al.* Heart failure with preserved, borderline, and reduced ejection fraction: 5-year outcomes. *J Am Coll Cardiol* 2017; 70:2476–2486.
- Lam CS, Donal E, Kraigher-Krainer E, Vasan RS. Epidemiology and clinical course of heart failure with preserved ejection fraction. *Eur J Heart Fail* 2011; 13:18–28.
- Kao DP, Lewsey JD, Anand IS, Massie BM, Zile MR, Carson PE, *et al.* Characterization of subgroups of heart failure patients with preserved ejection fraction with possible implications for prognosis and treatment response. *Eur J Heart Fail* 2015; 17:925–935.
- Ponikowski P, Voors AA, Anker SD, Bueno H, Cleland JGF, Coats AJS, *et al.* 2016 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure: The Task Force for the diagnosis and treatment of acute and chronic heart failure of the European Society of Cardiology (ESC) Developed with the special contribution of the Heart Failure Association (HFA) of the ESC. *Eur Heart J* 2016; 37:2129–2200.
- Yancy Clyde W, Jessup M, Bozkurt B, Butler J, Casey Donald E, Colvin Monica M, *et al.* 2017 ACC/AHA/HFSA focused update of the 2013 ACCF/AHA guideline for the management of heart failure: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines and the Heart Failure Society of America. *Circulation* 2017; 136:e137–e161.
- Okin PM, Devereux RB, Harris KE, Jern S, Kjeldsen SE, Julius S, *et al.* Regression of electrocardiographic left ventricular hypertrophy is associated with less hospitalization for heart failure in hypertensive patients. *Ann Intern Med* 2007; 147:311–319.
- Williams B, MacDonald TM, Morant S, Webb DJ, Sever P, McInnes G, *et al.* Spironolactone versus placebo, bisoprolol, and doxazosin to determine the optimal treatment for drug-resistant hypertension (PATHWAY-2): a randomised, double-blind, crossover trial. *Lancet* 2015; 386:2059–2068.
- Lam CSP, Voors AA, de Boer RA, Solomon SD, van Veldhuisen DJ. Heart failure with preserved ejection fraction: from mechanisms to therapies. *Eur Heart J* 2018; 39:2780–2792.
- Tsioufis C, Georgios G, Dimitrios O, Costas T, Niki K, Alexandros K, *et al.* Hypertension and heart failure with preserved ejection fraction: connecting the dots. *Curr Vasc Pharmacol* 2017; 16:15–22.
- Kjeldsen Sverre E, von Lueder Thomas G, Smiseth Otto A, Wachtell K, Mistry N, Westheim Arne S, *et al.* Medical therapies for heart failure with preserved ejection fraction. *Hypertension* 2020; 75:23–32.
- Jia G, Aroor Annayya R, Hill Michael A, Sowers James R. Role of renin-angiotensin-aldosterone system activation in promoting cardiovascular fibrosis and stiffness. *Hypertension* 2018; 72:537–548.
- Borlaug BA. The pathophysiology of heart failure with preserved ejection fraction. *Nat Rev Cardiol* 2014; 11:507–515.
- Senni M, Paulus WJ, Gavazzi A, Fraser AG, Diez J, Solomon SD, *et al.* New strategies for heart failure with preserved ejection fraction: the importance of targeted therapies for heart failure phenotypes. *Eur Heart J* 2014; 35:2797–2815.
- Shah SJ, Kitzman Dalane W, Borlaug Barry A, van Heerebeek L, Zile Michael R, Kass David A, *et al.* Phenotype-specific treatment of heart failure with preserved ejection fraction: a multiorgan roadmap. *Circulation* 2016; 134:73–90.
- Segar MW, Patel KV, Ayers C, Basit M, Tang WHW, Willett D, *et al.* Phenomapping of patients with heart failure with preserved ejection fraction using machine learning-based unsupervised cluster analysis. *Eur J Heart Fail* 2020; 22:148–158.
- Flint KM, Shah SJ, Lewis EF, Kao DP. Variation in clinical and patient-reported outcomes among complex heart failure with preserved ejection fraction phenotypes. *ESC Heart Fail* 2020; 7:811–824.
- Milani RV, Drazner MH, Lavie CJ, Morin DP, Ventura HO. Progression from concentric left ventricular hypertrophy and normal ejection fraction to left ventricular dysfunction. *Am J Cardiol* 2011; 108:992–996.
- Messerli FH, Rimoldi SF, Bangalore S. The transition from hypertension to heart failure: contemporary update. *JACC Heart Fail* 2017; 5:543–551.
- Pieske B, Tschöpe C, de Boer RA, Fraser AG, Anker SD, Donal E, *et al.* How to diagnose heart failure with preserved ejection fraction: the HFA-PEFF diagnostic algorithm: a consensus recommendation from the Heart Failure Association (HFA) of the European Society of Cardiology (ESC). *Eur Heart J* 2019; 40:3297–3317.
- Ho JE, Redfield Margaret M, Lewis Gregory D, Paulus Walter J, Lam Carolyn SP. Deliberating the diagnostic dilemma of heart failure with preserved ejection fraction. *Circulation* 2020; 142:1770–1780.
- Zile Michael R, Gottdiener John S, Hetzel Scott J, McMurray John J, Komajda M, McKelvie R, *et al.* Prevalence and significance of alterations in cardiac structure and function in patients with heart failure and a preserved ejection fraction. *Circulation* 2011; 124:2491–2501.
- Shah AM, Cikes M, Prasad N, Li G, Getchevski S, Claggett B, *et al.* Echocardiographic features of patients with heart failure and preserved left ventricular ejection fraction. *J Am Coll Cardiol* 2019; 74:2858–2873.
- Katz DH, Beussink L, Sauer AJ, Freed BH, Burke MA, Shah SJ. Prevalence, clinical characteristics, and outcomes associated with eccentric versus concentric left ventricular hypertrophy in heart failure with preserved ejection fraction. *Am J Cardiol* 2013; 112:1158–1164.
- Lam CS, Roger Véronique L, Rodeheffer Richard J, Bursi F, Borlaug Barry A, Ommen Steve R, *et al.* Cardiac structure and ventricular-vascular function in persons with heart failure and preserved ejection fraction from olmed county, minnesota. *Circulation* 2007; 115:1982–1990.
- Ganau A, Devereux RB, Roman MJ, de Simone G, Pickering TG, Saba PS, *et al.* Patterns of left ventricular hypertrophy and geometric remodeling in essential hypertension. *J Am Coll Cardiol* 1992; 19:1550–1558.
- Triposkiadis F, Butler J, Abboud FM, Armstrong PW, Adamopoulos S, Atherton JJ, *et al.* The continuous heart failure spectrum: moving beyond an ejection fraction classification. *Eur Heart J* 2019; 40:2155–2163.
- Halliday BP, Senior R, Pennell DJ. Assessing left ventricular systolic function: from ejection fraction to strain analysis. *Eur Heart J* 2020; 42:789–797.
- Aurigemma GP, Silver KH, Priest MA, Gaasch WH. Geometric changes allow normal ejection fraction despite depressed myocardial shortening in hypertensive left ventricular hypertrophy. *J Am Coll Cardiol* 1995; 26:195–202.
- Brucks S, Little WC, Chao T, Kitzman DW, Wesley-Farrington D, Gandhi S, *et al.* Contribution of left ventricular diastolic dysfunction to heart failure regardless of ejection fraction. *Am J Cardiol* 2005; 95:603–606.
- Konstam Marvin A, Abboud François M. Ejection fraction: misunderstood and overrated (changing the paradigm in categorizing heart failure). *Circulation* 2017; 135:717–719.

32. de Simone G, Devereux Richard B, Koren Michael J, Mensah George A, Casale Paul N, Laragh John H. Midwall left ventricular mechanics. An independent predictor of cardiovascular risk in arterial hypertension. *Circulation* 1996; 93:259–265.
33. Wachtell K, Palmieri V, Olsen Michael H, Gerdtts E, Papademetriou V, Nieminen Markku S, et al. Change in systolic left ventricular performance after 3 years of antihypertensive treatment: the Losartan Intervention for Endpoint (LIFE) Study. *Circulation* 2002; 106:227–232.
34. Wachtell K, Rokkedal J, Bella JN, Aalto T, Dahlöf B, Smith G, et al. Effect of electrocardiographic left ventricular hypertrophy on left ventricular systolic function in systemic hypertension (the LIFE study). Losartan Intervention For Endpoint. *Am J Cardiol* 2001; 87:54–60.
35. Okin PM, Wachtell K, Gerdtts E, Dahlöf B, Devereux RB. Relationship of left ventricular systolic function to persistence or development of electrocardiographic left ventricular hypertrophy in hypertensive patients: implications for the development of new heart failure. *J Hypertens* 2014; 32:2472–2478.
36. Stokke TM, Hasselberg NE, Smedsrud MK, Sarvari SI, Haugaa KH, Smiseth OA, et al. Geometry as a confounder when assessing ventricular systolic function: comparison between ejection fraction and strain. *J Am Coll Cardiol* 2017; 70:942–954.
37. Saito M, Khan F, Stoklosa T, Iannaccone A, Negishi K, Marwick TH. Prognostic implications of lv strain risk score in asymptomatic patients with hypertensive heart disease. *JACC Cardiovasc Imaging* 2016; 9:911–921.
38. Galderisi M, Esposito R, Schiano-Lomoriello V, Santoro A, Ippolito R, Schiattarella P, et al. Correlates of global area strain in native hypertensive patients: a three-dimensional speckle-tracking echocardiography study. *Eur Heart J Cardiovasc Imaging* 2012; 13:730–738.
39. Shah AM, Claggett B, Sweitzer Nancy K, Shah Sanjiv J, Anand Inder S, Liu L, et al. Prognostic importance of impaired systolic function in heart failure with preserved ejection fraction and the impact of spironolactone. *Circulation* 2015; 132:402–414.
40. Puwanant S, Priester TC, Mookadam F, Bruce CJ, Redfield MM, Chandrasekaran K. Right ventricular function in patients with preserved and reduced ejection fraction heart failure. *Eur J Echocardiogr* 2009; 10:733–737.
41. Gorter TM, van Veldhuisen DJ, Bauersachs J, Borlaug BA, Celutkiene J, Coats AJS, et al. Right heart dysfunction and failure in heart failure with preserved ejection fraction: mechanisms and management. Position statement on behalf of the Heart Failure Association of the European Society of Cardiology. *Eur J Heart Fail* 2018; 20:16–37.
42. Gorter TM, Obokata M, Reddy YNV, Melenovsky V, Borlaug BA. Exercise unmasks distinct pathophysiologic features in heart failure with preserved ejection fraction and pulmonary vascular disease. *Eur Heart J* 2018; 39:2825–2835.
43. Lam CS, Roger VL, Rodeheffer RJ, Borlaug BA, Enders FT, Redfield MM. Pulmonary hypertension in heart failure with preserved ejection fraction: a community-based study. *J Am Coll Cardiol* 2009; 53:1119–1126.
44. Gorter TM, van Veldhuisen DJ, Voors AA, Hummel YM, Lam CSP, Berger RMF, et al. Right ventricular-vascular coupling in heart failure with preserved ejection fraction and pre- vs. post-capillary pulmonary hypertension. *Eur Heart J Cardiovasc Imaging* 2018; 19:425–432.
45. Obokata M, Reddy YNV, Melenovsky V, Pislaru S, Borlaug BA. Deterioration in right ventricular structure and function over time in patients with heart failure and preserved ejection fraction. *Eur Heart J* 2019; 40:689–697.
46. Kawaguchi M, Hay I, Fetts B, Kass David A. Combined ventricular systolic and arterial stiffening in patients with heart failure and preserved ejection fraction: implications for systolic and diastolic reserve limitations. *Circulation* 2003; 107:714–720.
47. Ikonomidis I, Aboyans V, Blacher J, Brodmann M, Brutsaert DL, Chirinos JA, et al. The role of ventricular–arterial coupling in cardiac disease and heart failure: assessment, clinical implications and therapeutic interventions. A consensus document of the European Society of Cardiology Working Group on Aorta & Peripheral Vascular Diseases, European Association of Cardiovascular Imaging, and Heart Failure Association. *Eur J Heart Fail* 2019; 21:402–424.
48. Weber T, Chirinos JA. Pulsatile arterial haemodynamics in heart failure. *Eur Heart J* 2018; 39:3847–3854.
49. Borlaug BA, Kass DA. Ventricular–vascular interaction in heart failure. *Heart Fail Clin* 2008; 4:23–36.
50. Schwartzberg S, Redfield MM, From AM, Sorajja P, Nishimura RA, Borlaug BA. Effects of vasodilation in heart failure with preserved or reduced ejection fraction: implications of distinct pathophysiologies on response to therapy. *J Am Coll Cardiol* 2012; 59:442–451.
51. Borlaug Barry A, Melenovsky V, Russell Stuart D, Kessler K, Pacak K, Becker Lewis C, et al. Impaired chronotropic and vasodilator reserves limit exercise capacity in patients with heart failure and a preserved ejection fraction. *Circulation* 2006; 114:2138–2147.
52. Meyer M, LeWinter Martin M. Heart rate and heart failure with preserved ejection fraction: time to slow β -blocker use? *Circ Heart Fail* 2019; 12:e006213.
53. Tromp J, Shen L, Jhund PS, Anand IS, Carson PE, Desai AS, et al. Age-related characteristics and outcomes of patients with heart failure with preserved ejection fraction. *J Am Coll Cardiol* 2019; 74:601–612.
54. Oghina S, Bougouin W, Bézard M, Kharoubi M, Komajda M, Cohen-Solal A, et al. The impact of patients with cardiac amyloidosis in HFpEF trials. *JACC Heart Fail* 2021; 9:169–178.
55. Russo D, Musumeci MB, Volpe M. The neglected issue of cardiac amyloidosis in trials on heart failure with preserved ejection fraction in the elderly. *Eur J Heart Fail* 2020; 22:1740–1741.
56. Borlaug Barry A, Redfield Margaret M, Melenovsky V, Kane Garvan C, Karon Barry L, Jacobsen Steven J, et al. Longitudinal changes in left ventricular stiffness: a community-based study. *Circ Heart Fail* 2013; 6:944–952.
57. Shah AM, Claggett B, Kitzman D, Biering-Sørensen T, Jensen Jan S, Cheng S, et al. Contemporary assessment of left ventricular diastolic function in older adults: The Atherosclerosis Risk in Communities Study. *Circulation* 2017; 135:426–439.
58. Weber T, Wassertheurer S, Hametner B, Moebus S, Pundt N, Mahabadi AA, et al. Cross-sectional analysis of pulsatile hemodynamics across the adult life span: reference values, healthy and early vascular aging: the Heinz Nixdorf Recall and the MultiGeneration Study. *J Hypertens* 2019; 37:2404–2413.
59. Brouwers FP, de Boer RA, van der Harst P, Voors AA, Gansevoort RT, Bakker SJ, et al. Incidence and epidemiology of new onset heart failure with preserved vs. reduced ejection fraction in a community-based cohort: 11-year follow-up of PREVEND. *Eur Heart J* 2013; 34:1424–1431.
60. Damman K, Valente MAE, Voors AA, O'Connor CM, van Veldhuisen DJ, Hillege HL. Renal impairment, worsening renal function, and outcome in patients with heart failure: an updated meta-analysis. *Eur Heart J* 2014; 35:455–469.
61. Ter Maaten JM, Voors AA. Renal dysfunction in heart failure with a preserved ejection fraction: cause or consequence? *Eur J Heart Fail* 2016; 18:113–114.
62. Park M, Hsu C-y, Li Y, Mishra RK, Keane M, Rosas SE, et al. Associations between kidney function and subclinical cardiac abnormalities in CKD. *J Am Soc Nephrol* 2012; 23:1725–1734.
63. Unger ED, Dubin RF, Deo R, Daruwalla V, Friedman JL, Medina C, et al. Association of chronic kidney disease with abnormal cardiac mechanics and adverse outcomes in patients with heart failure and preserved ejection fraction. *Eur J Heart Fail* 2016; 18:103–112.
64. Zannad F, Rossignol P. Cardiorenal syndrome revisited. *Circulation* 2018; 138:929–944.
65. Yusuf S, Pfeffer MA, Swedberg K, Granger CB, Held P, McMurray JJV, et al. Effects of candesartan in patients with chronic heart failure and preserved left-ventricular ejection fraction: the CHARM-Preserved Trial. *The Lancet* 2003; 362:777–781.
66. Cleland JGF, Tendera M, Adamus J, Freemantle N, Polonski L, Taylor J, et al. The perindopril in elderly people with chronic heart failure (PEP-CHF) study. *Eur Heart J* 2006; 27:2338–2345.
67. Massie BM, Carson PE, McMurray JJ, Komajda M, McKelvie R, Zile MR, et al. Irbesartan in patients with heart failure and preserved ejection fraction. *N Engl J Med* 2008; 359:2456–2467.
68. Pitt B, Pfeffer MA, Assmann SF, Boineau R, Anand IS, Claggett B, et al. Spironolactone for heart failure with preserved ejection fraction. *N Engl J Med* 2014; 370:1383–1392.
69. Solomon SD, McMurray JJV, Anand IS, Ge J, Lam CSP, Maggioni AP, et al. Angiotensin–neprilysin inhibition in heart failure with preserved ejection fraction. *N Engl J Med* 2019; 381:1609–1620.

70. Kuno T, Ueyama H, Fujisaki T, Briasouli A, Takagi H, Briasoulis A. Meta-analysis evaluating the effects of renin-angiotensin-aldosterone system blockade on outcomes of heart failure with preserved ejection fraction. *Am J Cardiol* 2020; 125:1187–1193.
71. Salah HM, Fudim M, Al'Aref SJ, Khan MS, Almarzooq ZI, Devabhaktuni SR, *et al.* Meta-analysis of efficacy of sacubitril/valsartan in heart failure with preserved ejection fraction. *Am J Cardiol* 2021; 145:165–168.
72. Devereux Richard B, Dahlöf B, Gerds E, Boman K, Nieminen Markku S, Papademetriou V, *et al.* Regression of hypertensive left ventricular hypertrophy by losartan compared with atenolol: the Losartan Intervention for Endpoint Reduction in Hypertension (LIFE) trial. *Circulation* 2004; 110:1456–1462.
73. Lund LH, Benson L, Dahlström U, Edner M. Association between use of renin-angiotensin system antagonists and mortality in patients with heart failure and preserved ejection fraction. *JAMA* 2012; 308:2108–2117.
74. Khan MS, Fonarow GC, Khan H, Greene SJ, Anker SD, Gheorghiuade M, *et al.* Renin-angiotensin blockade in heart failure with preserved ejection fraction: a systematic review and meta-analysis. *ESC Heart Fail* 2017; 4:402–408.
75. Selvaraj S, Claggett B, Shah SJ, Anand I, Rouleau JL, Desai AS, *et al.* Systolic blood pressure and cardiovascular outcomes in heart failure with preserved ejection fraction: an analysis of the TOPCAT trial. *Eur J Heart Fail* 2018; 20:483–490.
76. Selvaraj S, Claggett BL, Böhm M, Anker SD, Vaduganathan M, Zannad F, *et al.* Systolic blood pressure in heart failure with preserved ejection fraction treated with sacubitril/valsartan. *J Am Coll Cardiol* 2020; 75:1644–1656.
77. Gupta AK, Poulter NR, Dobson J, Eldridge S, Cappuccio FP, Caulfield M, *et al.* Ethnic differences in blood pressure response to first and second-line antihypertensive therapies in patients randomized in the ASCOT trial. *Am J Hypertens* 2010; 23:1023–1030.
78. Ogedegbe G, Shah NR, Phillips C, Goldfeld K, Roy J, Guo Y, *et al.* Comparative effectiveness of angiotensin-converting enzyme inhibitor-based treatment on cardiovascular outcomes in hypertensive blacks versus whites. *J Am Coll Cardiol* 2015; 66:1224–1233.
79. Benedict CR, Johnstone DE, Weiner DH, Bourassa MG, Bittner V, Kay R, *et al.* Relation of neurohumoral activation to clinical variables and degree of ventricular dysfunction: a report from the registry of studies of left ventricular dysfunction. SOLVD Investigators. *J Am Coll Cardiol* 1994; 23:1410–1420.
80. Vergaro G, Aimo A, Prontera C, Ghionzoli N, Arzilli C, Zyw L, *et al.* Sympathetic and renin-angiotensin-aldosterone system activation in heart failure with preserved, mid-range and reduced ejection fraction. *Int J Cardiol* 2019; 296:91–97.
81. Catena C, Verheyen N, Pilz S, Kraigher-Krainer E, Tomaschitz A, Sechi Leonardo A, *et al.* Plasma aldosterone and left ventricular diastolic function in treatment-naïve patients with hypertension: tissue-doppler imaging study. *Hypertension* 2015; 65:1231–1237.
82. Gallo G, Tocci G, Fogacci F, Battistoni A, Rubattu S, Volpe M. Blockade of the neurohormonal systems in heart failure with preserved ejection fraction: a contemporary meta-analysis. *Int J Cardiol* 2020; 316:172–179.
83. McKelvie RS. The CHARM program: the effects of candesartan for the management of patients with chronic heart failure. *Expert Rev Cardiovasc Ther* 2009; 7:9–16.
84. Rogers JK, Pocock SJ, McMurray JJV, Granger CB, Michelson EL, Östergren J, *et al.* Analysing recurrent hospitalizations in heart failure: a review of statistical methodology, with application to CHARM-Preserved. *Eur J Heart Fail* 2014; 16:33–40.
85. Lund LH, Claggett B, Liu J, Lam CS, Jhund PS, Rosano GM, *et al.* Heart failure with mid-range ejection fraction in CHARM: characteristics, outcomes and effect of candesartan across the entire ejection fraction spectrum. *Eur J Heart Fail* 2018; 20:1230–1239.
86. Ferreira JP, Dewan P, Jhund PS, Lorenzo-Almorós A, Duarte K, Petrie MC, *et al.* Covariate adjusted reanalysis of the I-Preserve trial. *Clin Res Cardiol* 2020; 109:1358–1365.
87. Solomon SD, Rizkala AR, Gong J, Wang W, Anand IS, Ge J, *et al.* Angiotensin receptor neprilysin inhibition in heart failure with preserved ejection fraction: rationale and design of the PARAGON-HF trial. *JACC Heart Fail* 2017; 5:471–482.
88. Cunningham JW, Claggett BL, O'Meara E, Prescott MF, Pfeffer MA, Shah SJ, *et al.* Effect of sacubitril/valsartan on biomarkers of extracellular matrix regulation in patients with HFpEF. *J Am Coll Cardiol* 2020; 76:503–514.
89. Lyle MA, Iyer SR, Redfield MM, Reddy YNV, Felker GM, Cappola TP, *et al.* Circulating neprilysin in patients with heart failure and preserved ejection fraction. *JACC Heart Fail* 2020; 8:70–80.
90. Ruilope LM, Dukat A, Böhm M, Lacourcière Y, Gong J, Lefkowitz MP. Blood-pressure reduction with LCZ696, a novel dual-acting inhibitor of the angiotensin II receptor and neprilysin: a randomised, double-blind, placebo-controlled, active comparator study. *Lancet* 2010; 375:1255–1266.
91. Schmieder RE, Wagner F, Mayr M, Delles C, Ott C, Keicher C, *et al.* The effect of sacubitril/valsartan compared to olmesartan on cardiovascular remodelling in subjects with essential hypertension: the results of a randomized, double-blind, active-controlled study. *Eur Heart J* 2017; 38:3308–3317.
92. McMurray JJV, Packer M, Desai AS, Gong J, Lefkowitz MP, Rizkala AR, *et al.* Angiotensin-neprilysin inhibition versus enalapril in heart failure. *N Engl J Med* 2014; 371:993–1004.
93. Solomon SD, Zile M, Pieske B, Voors A, Shah A, Kraigher-Krainer E, *et al.* Prospective comparison of ARNI with ARB on management of heart failure with preserved ejection fraction (PARAMOUNT) investigators. The angiotensin receptor neprilysin inhibitor LCZ696 in heart failure with preserved ejection fraction: a phase 2 double-blind randomised controlled trial. *Lancet* 2012; 380:1387–1395.
94. US Food and Drug Administration briefing document. Cardiovascular and Renal Drugs Advisory Committee Meeting, December 15, 2020: NDA 207620: Sacubitril/Valsartan. <https://www.fda.gov/media/144377/download>. [Accessed 3 March 2021]
95. McMurray John JV, Jackson Alice M, Lam Carolyn SP, Redfield Margaret M, Anand Inder S, Ge J, *et al.* Effects of sacubitril-valsartan versus valsartan in women compared with men with heart failure and preserved ejection fraction: insights from PARAGON-HF. *Circulation* 2020; 141:338–351.
96. Solomon Scott D, Vaduganathan M, Brian LC, Packer M, Zile M, Swedberg K, *et al.* Sacubitril/valsartan across the spectrum of ejection fraction in heart failure. *Circulation* 2020; 141:352–361.
97. Vaduganathan M, Claggett BL, Desai AS, Anker SD, Perrone SV, Janssens S, *et al.* Prior heart failure hospitalization, clinical outcomes, and response to sacubitril/valsartan compared with valsartan in HFpEF. *J Am Coll Cardiol* 2020; 75:245–254.
98. Mc Causland Finnian R, Lefkowitz Martin P, Claggett B, Anavekar Nagesh S, Senni M, Gori M, *et al.* Angiotensin-neprilysin inhibition and renal outcomes in heart failure with preserved ejection fraction. *Circulation* 2020; 142:1236–1245.
99. Jering KS, Zannad F, Claggett B, Mc Causland FR, Ferreira JP, Desai A, *et al.* Cardiovascular and renal outcomes of mineralocorticoid receptor antagonist use in PARAGON-HF. *JACC Heart Fail* 2020; 9:13–24.
100. Vaduganathan M, Jhund PS, Claggett BL, Packer M, Widimský J, Seferovic P, *et al.* A putative placebo analysis of the effects of sacubitril/valsartan in heart failure across the full range of ejection fraction. *Eur Heart J* 2020; 41:2356–2362.
101. Wachter R, Shah SJ, Cowie MR, Szecsódy P, Shi V, Ibram G, *et al.* Angiotensin receptor neprilysin inhibition versus individualized RAAS blockade: design and rationale of the PARALLAX trial. *ESC Heart Fail* 2020; 7:856–864.
102. Nicholls M. Late-breaking clinical trials - ESC Congress 2020. *Eur Heart J* 2020; 41:4223–4227.
103. Zannad F, Alla F, Douset B, Perez A, Pitt B. Limitation of excessive extracellular matrix turnover may contribute to survival benefit of spironolactone therapy in patients with congestive heart failure: insights from the randomized aldactone evaluation study (RALES). Rales Investigators. *Circulation* 2000; 102:2700–2706.
104. Bauersachs J, Heck M, Fraccarollo D, Hildemann SK, Ertl G, Wehling M, *et al.* Addition of spironolactone to angiotensin-converting enzyme inhibition in heart failure improves endothelial vasomotor dysfunction: role of vascular superoxide anion formation and endothelial nitric oxide synthase expression. *J Am Coll Cardiol* 2002; 39:351–358.
105. Edwards NC, Steeds RP, Stewart PM, Ferro CJ, Townend JN. Effect of spironolactone on left ventricular mass and aortic stiffness in early-stage chronic kidney disease: a randomized controlled trial. *J Am Coll Cardiol* 2009; 54:505–512.

106. Edelmann F, Wachter R, Schmidt AG, Kraigher-Krainer E, Colantonio C, Kamke W, *et al.* Effect of spironolactone on diastolic function and exercise capacity in patients with heart failure with preserved ejection fraction: the Aldo-DHF randomized controlled trial. *JAMA* 2013; 309:781–791.
107. Pfeffer Marc A, Claggett B, Assmann Susan F, Boineau R, Anand Inder S, Clausell N, *et al.* Regional variation in patients and outcomes in the treatment of preserved cardiac function heart failure with an aldosterone antagonist (TOPCAT) trial. *Circulation* 2015; 131:34–42.
108. de Denuis S, O'Meara E, Desai AS, Claggett B, Lewis EF, Leclair G, *et al.* Spironolactone metabolites in TOPCAT — new insights into regional variation. *N Engl J Med* 2017; 376:1690–1692.
109. Girerd N, Ferreira JP, Rossignol P, Zannad F. A tentative interpretation of the TOPCAT trial based on randomized evidence from the brain natriuretic peptide stratum analysis. *Eur J Heart Fail* 2016; 18:1411–1414.
110. Solomon SD, Claggett B, Lewis EF, Desai A, Anand I, Sweitzer NK, *et al.* Influence of ejection fraction on outcomes and efficacy of spironolactone in patients with heart failure with preserved ejection fraction. *Eur Heart J* 2016; 37:455–462.
111. Böhm M, Perez A-C, Jhund PS, Reil JC, Komajda M, Zile MR, *et al.* Relationship between heart rate and mortality and morbidity in the irbesartan patients with heart failure and preserved systolic function trial (I-Preserve). *Eur J Heart Fail* 2014; 16:778–787.
112. Luchner A, Burnett JC, Jougasaki M, Hense H-W, Riegger GAJ, Schunkert H. Augmentation of the cardiac natriuretic peptides by beta-receptor antagonism: evidence from a population-based study. *J Am Coll Cardiol* 1998; 32:1839–1844.
113. Pucci G, Ranalli Maria G, Battista F, Schillaci G. Effects of β -blockers with and without vasodilating properties on central blood pressure: systematic review and meta-analysis of randomized trials in hypertension. *Hypertension* 2016; 67:316–324.
114. Hernandez AF, Hammill BG, O'Connor CM, Schulman KA, Curtis LH, Fonarow GC. Clinical effectiveness of beta-blockers in heart failure: findings from the OPTIMIZE-HF (Organized Program to Initiate Lifesaving Treatment in Hospitalized Patients With Heart Failure) registry. *J Am Coll Cardiol* 2009; 53:184–192.
115. Yamamoto K, Origasa H, Hori M. Effects of carvedilol on heart failure with preserved ejection fraction: the Japanese Diastolic Heart Failure Study (J-DHF). *Eur J Heart Fail* 2013; 15:110–118; on behalf of the JDHFI.
116. Bavishi C, Chatterjee S, Ather S, Patel D, Messerli FH. Beta-blockers in heart failure with preserved ejection fraction: a meta-analysis. *Heart Fail Rev* 2015; 20:193–201.
117. Flather MD, Shibata MC, Coats AJS, Van Veldhuisen DJ, Parkhomenko A, Borbola J, *et al.* Randomized trial to determine the effect of nebivolol on mortality and cardiovascular hospital admission in elderly patients with heart failure (SENIORS). *Eur Heart J* 2005; 26:215–225.
118. Conraads VM, Metra M, Kamp O, De Keulenaer GW, Pieske B, Zamorano J, *et al.* Effects of the long-term administration of nebivolol on the clinical symptoms, exercise capacity, and left ventricular function of patients with diastolic dysfunction: results of the ELANDD study. *Eur J Heart Fail* 2012; 14:219–225.
119. Komajda M, Isnard R, Cohen-Solal A, Metra M, Pieske B, Ponikowski P, *et al.* Effect of ivabradine in patients with heart failure with preserved ejection fraction: the EDIFY randomized placebo-controlled trial. *Eur J Heart Fail* 2017; 19:1495–1503.
120. Cleland JGF, Bunting KV, Flather MD, Altman DG, Holmes J, Coats AJS, *et al.* Beta-blockers for heart failure with reduced, mid-range, and preserved ejection fraction: an individual patient-level analysis of double-blind randomized trials. *Eur Heart J* 2018; 39:26–35.
121. Silverman DN, Plante TB, Infeld M, Callas PW, Juraschek SP, Dougherty GB, *et al.* Association of (-)blocker use with heart failure hospitalizations and cardiovascular disease mortality among patients with heart failure with a preserved ejection fraction: a secondary analysis of the TOPCAT trial. *JAMA Netw Open* 2019; 2:e1916598.
122. Lam CS, Chandramouli C, Ahojja V, Verma S. SGLT-2 inhibitors in heart failure: current management, unmet needs, and therapeutic prospects. *J Am Heart Assoc* 2019; 8:e013389.
123. Buse JB, Wexler DJ, Tsapas A, Rossing P, Mingrone G, Mathieu C, *et al.* 2019 update to: management of hyperglycemia in type 2 diabetes, 2018. A Consensus Report by the American Diabetes Association (ADA) and the European Association for the Study of Diabetes (EASD). *Diabetes Care* 2020; 43:487.
124. Marx N, Davies MJ, Grant PJ, Mathieu C, Petrie JR, Cosentino F, *et al.* Guideline recommendations and the positioning of newer drugs in type 2 diabetes care. *Lancet Diabetes Endocrinol* 2020; 9:46–52.
125. Boyer JK, Thanigaraj S, Schechtman KB, Pérez JE. Prevalence of ventricular diastolic dysfunction in asymptomatic, normotensive patients with diabetes mellitus. *Am J Cardiol* 2004; 93:870–875.
126. Stahrenberg R, Edelmann F, Mende M, Kocksämper A, Dungen HD, Scherer M, *et al.* Association of glucose metabolism with diastolic function along the diabetic continuum. *Diabetologia* 2010; 53:1331–1340.
127. Aguilar D, Deswal A, Ramasubbu K, Mann DL, Bozkurt B. Comparison of patients with heart failure and preserved left ventricular ejection fraction among those with versus without diabetes mellitus. *Am J Cardiol* 2010; 105:373–377.
128. Cosentino F, Grant PJ, Aboyans V, Bailey CJ, Ceriello A, Delgado V, *et al.* 2019 ESC Guidelines on diabetes, pre-diabetes, and cardiovascular diseases developed in collaboration with the EASD: the Task Force for diabetes, pre-diabetes, and cardiovascular diseases of the European Society of Cardiology (ESC) and the European Association for the Study of Diabetes (EASD). *Eur Heart J* 2020; 41:255–323.
129. Chin KL, Ofori-Asenso R, Hopper I, von Lueder TG, Reid CM, Zoungas S, *et al.* Potential mechanisms underlying the cardiovascular benefits of sodium glucose cotransporter 2 inhibitors: a systematic review of data from preclinical studies. *Cardiovasc Res* 2019; 115:266–276.
130. Hallow KM, Helmlinger G, Greasley PJ, McMurray JVV, Boulton DW. Why do SGLT2 inhibitors reduce heart failure hospitalization? A differential volume regulation hypothesis. *Diabetes Obes Metab* 2018; 20:479–487.
131. Striepe K, Jumar A, Ott C, Karg Marina V, Schneider Markus P, Kannenkeril D, *et al.* Effects of the selective sodium-glucose cotransporter 2 inhibitor empagliflozin on vascular function and central hemodynamics in patients with type 2 diabetes mellitus. *Circulation* 2017; 136:1167–1169.
132. Verma S, Mazer CD, Yan Andrew T, Mason T, Garg V, Teoh H, *et al.* Effect of empagliflozin on left ventricular mass in patients with type 2 diabetes mellitus and coronary artery disease: the EMPA-HEART CardioLink-6 randomized clinical trial. *Circulation* 2019; 140:1693–1702.
133. Pabel S, Wagner S, Bollenberg H, Bengel P, Kovács Á, Schach C, *et al.* Empagliflozin directly improves diastolic function in human heart failure. *Eur J Heart Fail* 2018; 20:1690–1700.
134. Verma S, Rawat S, Ho KL, Wagg CS, Zhang L, Teoh H, *et al.* Empagliflozin increases cardiac energy production in diabetes: novel translational insights into the heart failure benefits of SGLT2 inhibitors. *JACC Basic Transl Sci* 2018; 3:575–587.
135. Zinman B, Wanner C, Lachin JM, Fitchett D, Bluhmki E, Hantel S, *et al.* Empagliflozin, cardiovascular outcomes, and mortality in type 2 diabetes. *N Engl J Med* 2015; 373:2117–2128.
136. Perkovic V, Jardine MJ, Neal B, Bompoint S, Heerspink HJL, Charytan DM, *et al.* Canagliflozin and renal outcomes in type 2 diabetes and nephropathy. *N Engl J Med* 2019; 380:2295–2306.
137. Wiviott SD, Raz I, Bonaca MP, Mosenzon O, Kato ET, Cahn A, *et al.* Dapagliflozin and cardiovascular outcomes in type 2 diabetes. *N Engl J Med* 2019; 380:347–357.
138. Cannon CP, Pratley R, Dagogo-Jack S, Mancuso J, Huyck S, Masiukiewicz U, *et al.* Cardiovascular outcomes with ertugliflozin in type 2 diabetes. *N Engl J Med* 2020; 383:1425–1435.
139. McMurray JVV, Solomon SD, Inzucchi SE, Køber L, Kosiborod MN, Martinez FA, *et al.* Dapagliflozin in patients with heart failure and reduced ejection fraction. *N Engl J Med* 2019; 381:1995–2008.
140. Packer M, Anker SD, Butler J, Filippatos G, Pocock SJ, Carson P, *et al.* Cardiovascular and renal outcomes with empagliflozin in heart failure. *N Engl J Med* 2020; 383:1413–1424.
141. Zannad F, Ferreira JP, Pocock SJ, Anker SD, Butler J, Filippatos G, *et al.* SGLT2 inhibitors in patients with heart failure with reduced ejection fraction: a meta-analysis of the EMPEROR-Reduced and DAPA-HF trials. *Lancet* 2020; 396:819–829.
142. Seferović PM, Coats AJS, Ponikowski P, Filippatos G, Huelsmann M, Jhund PS, *et al.* European Society of Cardiology/Heart Failure Association position paper on the role and safety of new glucose-lowering drugs in patients with heart failure. *Eur J Heart Fail* 2020; 22:196–213.

143. Seferović PM, Fragasso G, Petrie M, Mullens W, Ferrari R, Thum T, *et al.* Sodium–glucose co-transporter 2 inhibitors in heart failure: beyond glycaemic control. A position paper of the Heart Failure Association of the European Society of Cardiology. *Eur J Heart Fail* 2020; 22:1495–1503.
144. Seferović PM, Fragasso G, Petrie M, Mullens W, Ferrari R, Thum T, *et al.* Heart Failure Association of the European Society of Cardiology update on sodium–glucose co-transporter 2 inhibitors in heart failure. *Eur J Heart Fail* 2020; 22:1984–1986.
145. Januzzi J, Ferreira JP, Böhm M, Kaul S, Wanner C, Brueckmann M, *et al.* Empagliflozin reduces the risk of a broad spectrum of heart failure outcomes regardless of heart failure status at baseline. *Eur J Heart Fail* 2019; 21:386–388.
146. Bhatt DL, Szarek M, Steg PG, Cannon CP, Leiter LA, McGuire DK, *et al.* Sotagliflozin in patients with diabetes and recent worsening heart failure. *N Engl J Med* 2020; 384:117–128.
147. Volpe M, Patrono C. The value of sotagliflozin in patients with diabetes and heart failure detracted by an unexpected ending. *Eur Heart J* 2021; 42:1458–1459.
148. Tschöpe C, Birner C, Böhm M, Bruder O, Frantz S, Luchner A, *et al.* Heart failure with preserved ejection fraction: current management and future strategies: expert opinion on the behalf of the Nucleus of the “Heart Failure Working Group” of the German Society of Cardiology (DKG). *Clin Res Cardiol* 2018; 107:1–19.
149. Wintrich J, Kindermann I, Ukena C, Selejan S, Werner C, Maack C, *et al.* Therapeutic approaches in heart failure with preserved ejection fraction: past, present, and future. *Clin Res Cardiol* 2020; 109:1079–1098.
150. Redfield MM, Chen HH, Borlaug BA, Semigran MJ, Lee KL, Lewis G, *et al.* Effect of phosphodiesterase-5 inhibition on exercise capacity and clinical status in heart failure with preserved ejection fraction: a randomized clinical trial. *JAMA* 2013; 309:1268–1277.
151. Shah SJ, Feldman T, Ricciardi MJ, Kahwash R, Lilly S, Litwin S, *et al.* One-year safety and clinical outcomes of a transcatheter interatrial shunt device for the treatment of heart failure with preserved ejection fraction in the Reduce Elevated Left Atrial Pressure in Patients With Heart Failure (REDUCE LAP-HF I) trial: a randomized clinical trial. *JAMA Cardiol* 2018; 3:968–977.
152. Tsioufis C, Papademetriou V, Dimitriadis K, Tsiachris D, Thomopoulos C, Kasiakogias A, *et al.* Effects of multielectrode renal denervation on cardiac and neurohumoral adaptations in resistant hypertension with cardiac hypertrophy: an EnligHTN I substudy. *J Hypertens* 2015; 33:346–353.
153. Kandzari DE, Böhm M, Mahfoud F, Townsend RR, Weber MA, Pocock S, *et al.* Effect of renal denervation on blood pressure in the presence of antihypertensive drugs: 6-month efficacy and safety results from the SPYRAL HTN-ON MED proof-of-concept randomised trial. *Lancet* 2018; 391:2346–2355.
154. Böhm M, Kario K, Kandzari DE, Mahfoud F, Weber MA, Schmieder RE, *et al.* Efficacy of catheter-based renal denervation in the absence of antihypertensive medications (SPYRAL HTN-OFF MED Pivotal): a multicentre, randomised, sham-controlled trial. *Lancet* 2020; 395:1444–1451.
155. Patel HC, Rosen SD, Hayward C, Vassiliou V, Smith GC, Wage RR, *et al.* Renal denervation in heart failure with preserved ejection fraction (RDT-PEF): a randomized controlled trial. *Eur J Heart Fail* 2016; 18:703–712.
156. Ather S, Chan W, Bozkurt B, Aguilar D, Ramasubbu K, Zachariah AA, *et al.* Impact of noncardiac comorbidities on morbidity and mortality in a predominantly male population with heart failure and preserved versus reduced ejection fraction. *J Am Coll Cardiol* 2012; 59:998–1005.
157. Camafort M, Kario K. Hypertension, heart failure, and frailty in older people: a common but unclear situation. *J Clin Hypertens* 2020; 22:1763–1768.
158. Devereux RB, Agabiti-Rosei E, Dahlöf B, Gosse P, Hahn RT, Okin PM, *et al.* Regression of left ventricular hypertrophy as a surrogate endpoint for morbid events in hypertension treatment trials. *J Hypertens* 1996; 14:S95–101.
159. Fagard Robert H, Celis H, Thijs L, Wouters S. Regression of left ventricular mass by antihypertensive treatment: a meta-analysis of randomized comparative studies. *Hypertension* 2009; 54:1084–1091.
160. Jekell A, Nilsson PM, Kahan T. Treatment of hypertensive left ventricular hypertrophy. *Curr Pharm Design* 2018; 24:4391–4396.
161. Salvetti M, Paini A, Bertacchini F, Stassaldi D, Aggiusti C, Agabiti Rosei C, *et al.* Changes in left ventricular geometry during antihypertensive treatment. *Pharmacol Res* 2018; 134:193–199.
162. Wachtell K, Bella Jonathan N, Rokkedal J, Palmieri V, Papademetriou V, Dahlöf B, *et al.* Change in diastolic left ventricular filling after one year of antihypertensive treatment: the losartan intervention for endpoint reduction in hypertension (LIFE) study. *Circulation* 2002; 105:1071–1076.
163. Okin PM, Devereux RB, Jern S, Kjeldsen SE, Julius S, Nieminen MS, *et al.* Regression of electrocardiographic left ventricular hypertrophy during antihypertensive treatment and the prediction of major cardiovascular events. *JAMA* 2004; 292:2343–2349.
164. Kawano H, Fujiwara A, Kai H, Kumagai E, Okamoto R, Shibata R, *et al.* Effects of blood pressure lowering in patients with heart failure with preserved ejection fraction: a systematic review and meta-analysis. *Hypertens Res* 2019; 42:504–513.
165. Upadhyaya B, Rocco M, Lewis Cora E, Oparil S, Lovato Laura C, Cushman William C, *et al.* Effect of intensive blood pressure treatment on heart failure events in the systolic blood pressure reduction intervention trial. *Circ Heart Fail* 2017; 10:e003613.
166. Kostis JB, Davis BR, Cutler J, Grimm RH Jr, Berge KG, Cohen JD, *et al.* Prevention of heart failure by antihypertensive drug treatment in older persons with isolated systolic hypertension. SHEP Cooperative Research Group. *JAMA* 1997; 278:212–216.
167. Thomopoulos C, Parati G, Zanchetti A. Effects of blood pressure-lowering treatment. 6. Prevention of heart failure and new-onset heart failure – meta-analyses of randomized trials. *J Hypertens* 2016; 34:373–384.
168. Huang P, Yu Y, Wei F, Zhu W, Xue R, Dong Y, *et al.* Association of long-term SBP with clinical outcomes and quality of life in heart failure with preserved ejection fraction: an analysis of the Treatment of Preserved Cardiac Function Heart Failure with an Aldosterone Antagonist trial. *J Hypertens* 2021 [Epub ahead of print].
169. Rossignol P, Claggett BL, Liu J, Vardeny O, Pitt B, Zannad F, *et al.* Spironolactone and resistant hypertension in heart failure with preserved ejection fraction. *Am J Hypertens* 2018; 31:407–414.
170. Faselis C, Lam PH, Zile MR, Bhyan P, Tsimploulis A, Arundel C, *et al.* Systolic blood pressure and outcomes in older patients with HFpEF and hypertension. *Am J Med* 2021; 134:e252–e263.
171. Tsimploulis A, Lam PH, Arundel C, Singh SN, Morgan CJ, Faselis C, *et al.* Systolic blood pressure and outcomes in patients with heart failure with preserved ejection fraction. *JAMA Cardiol* 2018; 3:288–297.
172. Komori T, Eguchi K, Saito T, Hoshida S, Kario K. Riser pattern is a novel predictor of adverse events in heart failure patients with preserved ejection fraction. *Circ J* 2017; 81:220–226.
173. Ueda T, Kawakami R, Nakada Y, Nakano T, Nakagawa H, Matsui M, *et al.* Differences in blood pressure riser pattern in patients with acute heart failure with reduced mid-range and preserved ejection fraction. *ESC Heart Fail* 2019; 6:1057–1067.
174. Lee SE, Lee H-Y, Cho H-J, Choe W-S, Kim H, Choi J-O, *et al.* Reverse j-curve relationship between on-treatment blood pressure and mortality in patients with heart failure. *JACC Heart Fail* 2017; 5:810–819.
175. Tadic M, Cuspidi C. Obesity and heart failure with preserved ejection fraction: a paradox or something else? *Heart Fail Rev* 2019; 24:379–385.
176. Obokata M, Reddy Yogesh NV, Pislaru Sorin V, Melenovsky V, Borlaug Barry A. Evidence supporting the existence of a distinct obese phenotype of heart failure with preserved ejection fraction. *Circulation* 2017; 136:6–19.
177. Kitzman DW, Brubaker P, Morgan T, Haykowsky M, Hundley G, Kraus WE, *et al.* Effect of caloric restriction or aerobic exercise training on peak oxygen consumption and quality of life in obese older patients with heart failure with preserved ejection fraction: a randomized clinical trial. *JAMA* 2016; 315:36–46.
178. Khan MS, Jones DW, Butler J. Salt, no salt, or less salt for patients with heart failure? *Am J Med* 2020; 133:32–38.
179. Hummel Scott L, Seymour EM, Brook Robert D, Sheth Samar S, Ghosh E, Zhu S, *et al.* Low-sodium dash diet improves diastolic function and ventricular–arterial coupling in hypertensive heart failure with preserved ejection fraction. *Circ Heart Fail* 2013; 6:1165–1171.

180. Edelmann F, Gelbrich G, Dungen H-D, Fröhling S, Wachter R, Stahrenberg R, *et al.* Exercise training improves exercise capacity and diastolic function in patients with heart failure with preserved ejection fraction: results of the Ex-DHF (Exercise training in Diastolic Heart Failure) pilot study. *J Am Coll Cardiol* 2011; 58:1780–1791.
181. Hegde Sheila M, Claggett B, Shah Amil M, Lewis Eldrin F, Anand I, Shah Sanjiv J, *et al.* Physical activity and prognosis in the TOPCAT trial (treatment of preserved cardiac function heart failure with an aldosterone antagonist). *Circulation* 2017; 136:982–992.
182. Fukuta H, Goto T, Wakami K, Kamiya T, Ohte N. Effects of exercise training on cardiac function, exercise capacity, and quality of life in heart failure with preserved ejection fraction: a meta-analysis of randomized controlled trials. *Heart Fail Rev* 2019; 24:535–547.
183. Mueller S, Winzer EB, Duvinage A, Gevaert AB, Edelmann F, Haller B, *et al.* Effect of high-intensity interval training, moderate continuous training, or guideline-based physical activity advice on peak oxygen consumption in patients with heart failure with preserved ejection fraction: a randomized clinical trial. *JAMA* 2021; 325:542–551.
184. PellICCIA A, Sharma S, Gati S, Bäck M, Börjesson M, Caselli S, *et al.* 2020 ESC Guidelines on sports cardiology and exercise in patients with cardiovascular disease: the Task Force on sports cardiology and exercise in patients with cardiovascular disease of the European Society of Cardiology (ESC). *Eur Heart J* 2021; 42:17–96.
185. Laoutaris ID, Piotrowicz E, Kallistratos MS, Dritsas A, Dimaki N, Miliopoulos D, *et al.* Combined aerobic/resistance/inspiratory muscle training as the ‘optimum’ exercise programme for patients with chronic heart failure: ARISTOS-HF randomized clinical trial. *Eur J Prev Cardiol* 2021.
186. Warner James G, Metzger DC, Kitzman Dalane W, Wesley Deborah J, Little William C. Losartan improves exercise tolerance in patients with diastolic dysfunction and a hypertensive response to exercise. *J Am Coll Cardiol* 1999; 33:1567–1572.
187. Solomon SD, Janardhanan R, Verma A, Bourgoun M, Daley WL, Purkayastha D, *et al.* Effect of angiotensin receptor blockade and antihypertensive drugs on diastolic function in patients with hypertension and diastolic dysfunction: a randomised trial. *Lancet* 2007; 369:2079–2087.
188. Faselis C, Arundel C, Patel S, Lam PH, Gottlieb SS, Zile MR, *et al.* Loop diuretic prescription and 30-day outcomes in older patients with heart failure. *J Am Coll Cardiol* 2020; 76:669–679.
189. Mullens W, Damman K, Harjola VP, Mebazaa A, Brunner-La Rocca HP, Martens P, *et al.* The use of diuretics in heart failure with congestion: a position statement from the Heart Failure Association of the European Society of Cardiology. *Eur J Heart Fail* 2019; 21:137–155.
190. Costanzo MR, Stevenson LW, Adamson PB, Desai AS, Heywood JT, Bourge RC, *et al.* Interventions linked to decreased heart failure hospitalizations during ambulatory pulmonary artery pressure monitoring. *JACC Heart Fail* 2016; 4:333–344.
191. Adamson Philip B, Abraham William T, Bourge Robert C, Costanzo Maria R, Hasan A, Yadav C, *et al.* Wireless pulmonary artery pressure monitoring guides management to reduce decompensation in heart failure with preserved ejection fraction. *Circ Heart Fail* 2014; 7:935–944.
192. López B, Querejeta R, González A, Beaumont J, Larman M, Díez J. Impact of treatment on myocardial lysyl oxidase expression and collagen cross-linking in patients with heart failure. *Hypertension* 2009; 53:236–242.
193. US Food and Drug Administration briefing document. Cardiovascular and Renal Drugs Advisory Committee Meeting, December 16, 2020: Spironolactone for Heart Failure with Preserved Ejection Fraction (HFpEF). <https://www.fda.gov/media/144411/download>.
194. US Food and Drug Administration Center for Drug Evaluation and Research. Final Summary Minutes of the Cardiovascular and Renal Drugs Advisory Committee Meeting, December 16, 2020. <https://www.fda.gov/media/145548/download>. [Accessed 3 March 2021]
195. Ravassa S, Trippel T, Bach D, Bachran D, González A, López B, *et al.* Biomarker-based phenotyping of myocardial fibrosis identifies patients with heart failure with preserved ejection fraction resistant to the beneficial effects of spironolactone: results from the Aldo-DHF trial. *Eur J Heart Fail* 2018; 20:1290–1299.
196. Cleland JGF, Ferreira JP, Mariottoni B, Pellicori P, Cuthbert J, Verdonschot JAJ, *et al.* The effect of spironolactone on cardiovascular function and markers of fibrosis in people at increased risk of developing heart failure: the heart ‘OMics’ in AGEing (HOMAGE) randomized clinical trial. *Eur Heart J* 2020; 42:684–696.
197. Müller-Brunotte R, Kahan T, López B, Edner M, González A, Díez J, *et al.* Myocardial fibrosis and diastolic dysfunction in patients with hypertension: results from the Swedish Irbesartan Left Ventricular Hypertrophy Investigation versus Atenolol (SILVHIA). *J Hypertens* 2007; 25:1958–1966.
198. González A, López B, Querejeta R, Zubillaga E, Echeverría T, Díez J. Filling pressures and collagen metabolism in hypertensive patients with heart failure and normal ejection fraction. *Hypertension* 2010; 55:1418–1424.
199. Suzuki K, Claggett B, Minamisawa M, Nochioka K, Mitchell GF, Anand IS, *et al.* Pulse pressure, prognosis, and influence of sacubitril/valsartan in heart failure with preserved ejection fraction. *Hypertension* 2021; 77:546–556.
200. US Food and Drug Administration Center for Drug Evaluation and Research. Final Summary Minutes of the Cardiovascular and Renal Drugs Advisory Committee Meeting, December 15, 2020. <https://www.fda.gov/media/145549/download>. [Accessed 3 March 2021]
201. ENTRESTO [prescribing information]; East Hanover, NJ: Novartis Pharmaceuticals Corp. <https://www.novartis.us/sites/www.novartis.us/files/entresto.pdf>. [Accessed 3 March 2021]
202. Setaro JF, Zaret BL, Schulman DS, Black HR, Soufer R. Usefulness of verapamil for congestive heart failure associated with abnormal left ventricular diastolic filling and normal left ventricular systolic performance. *Am J Cardiol* 1990; 66:981–986.
203. Packer M, Carson P, Elkayam U, Konstam MA, Moe G, O’Connor C, *et al.* Effect of amlodipine on the survival of patients with severe chronic heart failure due to a nonischemic cardiomyopathy: results of the PRAISE-2 study (prospective randomized amlodipine survival evaluation 2). *JACC Heart Fail* 2013; 1:308–314.
204. Patel K, Fonarow Gregg C, Ahmed M, Morgan C, Kilgore M, Love Thomas E, *et al.* Calcium channel blockers and outcomes in older patients with heart failure and preserved ejection fraction. *Circ Heart Fail* 2014; 7:945–952.
205. Thomopoulos C, Bazoukis G, Tsioufis C, Mancia G. Beta-blockers in hypertension: overview and meta-analysis of randomized outcome trials. *J Hypertens* 2020; 38:1669–1681.
206. Mancia G, Masi S, Palatini P, Tsioufis C, Grassi G. Elevated heart rate and cardiovascular risk in hypertension. *J Hypertens* 2021. [Epub ahead of print].
207. Schelbert EB, Fridman Y, Wong TC, Abu Daya H, Piehler KM, Kadakkal A, *et al.* Temporal relation between myocardial fibrosis and heart failure with preserved ejection fraction: association with baseline disease severity and subsequent outcome. *JAMA Cardiol* 2017; 2:995–1006.
208. Díez J, Querejeta R, López B, González A, Larman M, Martínez Ubago José L. losartan-dependent regression of myocardial fibrosis is associated with reduction of left ventricular chamber stiffness in hypertensive patients. *Circulation* 2002; 105:2512–2517.
209. McDiarmid Adam K, Swoboda Peter P, Erhayiem B, Bounford Katrina A, Bijsterveld P, Tyndall K, *et al.* Myocardial effects of aldosterone antagonism in heart failure with preserved ejection fraction. *J Am Heart Assoc* 2020; 9:e011521.
210. Reddy YNV, Obokata M, Verbrugge FH, Lin G, Borlaug BA. Atrial dysfunction in patients with heart failure with preserved ejection fraction and atrial fibrillation. *J Am Coll Cardiol* 2020; 76:1051–1064.
211. Katz DH, Deo RC, Aguilar FG, Selvaraj S, Martinez EE, Beussink-Nelson L, *et al.* Phenomapping for the identification of hypertensive patients with the myocardial substrate for heart failure with preserved ejection fraction. *J Cardiovasc Transl Res* 2017; 10:275–284.
212. Smiseth OA, Torp H, Opdahl A, Haugaa KH, Urheim S. Myocardial strain imaging: how useful is it in clinical decision making? *Eur Heart J* 2016; 37:1196–1207.
213. Gu H, Li Y, Fok H, Simpson J, Kentish Jonathan C, Shah Ajay M, *et al.* Reduced first-phase ejection fraction and sustained myocardial wall stress in hypertensive patients with diastolic dysfunction: a manifestation of impaired shortening deactivation that links systolic to diastolic dysfunction and preserves systolic ejection fraction. *Hypertension* 2017; 69:633–640.

214. Cleland John G, Pellicori P, Januzzi J, Zannad F, Clark AL, McMurray John J, *et al.* The conceptual basis for a Universal Definition of Heart Failure: congestion due to cardiac dysfunction. *Eur Heart J* 2021;0.
215. Bauersachs J, Cleland John G, Pellicori P, Januzzi J, Zannad F, Clark AL, *et al.* The struggle towards a Universal Definition of Heart Failure: how to proceed? *Eur Heart J* 2021; 0:1–13.
216. Pfeffer Marc A, Teerlink JR. Counter point: no evidence for the proposed Universal Definition of Heart Failure. *Eur Heart J* 2021; 0:1–13.
217. Abraham WT, Psotka MA, Fiuzat M, Filippatos G, Lindenfeld J, Mehran R, *et al.* Standardized definitions for evaluation of heart failure therapies: scientific expert panel from the Heart Failure Collaboratory and Academic Research Consortium. *JACC Heart Fail* 2020; 8:961–972.
218. Anand Inder S, Rector Thomas S, Cleland John G, Kuskowski M, McKelvie Robert S, Persson H, *et al.* Prognostic value of baseline plasma amino-terminal pro-brain natriuretic peptide and its interactions with irbesartan treatment effects in patients with heart failure and preserved ejection fraction: findings from the I-PRESERVE trial. *Circ Heart Fail* 2011; 4:569–577.
219. Anand IS, Claggett B, Liu J, Shah AM, Rector TS, Shah SJ, *et al.* Interaction between spironolactone and natriuretic peptides in patients with heart failure and preserved ejection fraction: from the TOPCAT Trial. *JACC Heart Fail* 2017; 5:241–252.
220. Cunningham JW, Vaduganathan M, Claggett BL, Zile MR, Anand IS, Packer M, *et al.* Effects of sacubitril/valsartan on N-Terminal Pro-B-Type Natriuretic Peptide in heart failure with preserved ejection fraction. *JACC Heart Fail* 2020; 8:372–381.